

# Chemistry 1A Fall 2000

## Midterm Exam II, version A October 17, 2000

(Closed book, 75 minutes, 145 points)

Name: \_\_\_\_\_

Section Number: \_\_\_\_\_

SID: \_\_\_\_\_

T.A. Name: \_\_\_\_\_

Identification Sticker

Exam information, extra directions, and useful hints to maximize your score:

- Write your name on all ten pages.
- There are two parts to the exam: 1) multiple choice and 2) short answer problems.
- **For the multiple choice problems, fill in the Scantron™ form AND circle the answer on your exam.**
- Answer the questions you know how to do first, then work on the questions you skipped.
- Show all work for which you want credit on the short answer problems and do not forget units!
- You may use the back side of the exam pages for scratch paper.

### Unit Prefixes

milli, m ( $\times 10^{-3}$ )	micro, $\mu$ ( $\times 10^{-6}$ )	nano, n ( $\times 10^{-9}$ )
kilo, k ( $\times 10^3$ )	mega, M ( $\times 10^6$ )	giga, G ( $\times 10^9$ )

Some possibly useful information:

$$E_{\text{photon}} = h\nu = hc/\lambda$$

$$A = \epsilon c$$

$$v_{\text{rms}} = \sqrt{\frac{3RT}{M}}$$

$$E_{\text{kin}} = \frac{3}{2} nRT$$


$$PV = nRT$$

$$E_n = -\frac{Z^2}{n^2} R_{\infty}$$

	Ionization Energy (kJ/mol)	Electron Affinity (kJ/mol)
Na	496	-53
Ne	2081	~0

$$T(\text{K}) = T(^{\circ}\text{C}) + 273.15$$

(Do not write in this box; it is for official use only.)






Page	Points
2-5	/ 60
6-7	/ 40
8-10	/ 45
Total	/ 145

**Part 1: Multiple Choice (5 pts each, 60 pts total)**


Instructions: Bubble in the correct answer on your Scantron sheet AND circle the answer on your exam. Each question has one correct answer.

- 1.) The answer to question 1 is **A**. Bubble in **A** on your Scantron™ form.
- 2.) What is the electron affinity of  $\text{Na}^+$ ?
- A.) -2081 kJ/mol                      B.) -496 kJ/mol                      C.) ~0 kJ/mol
- D.) 53 kJ/mol                              E.) 2081 kJ/mol


- 3.) Select the correct energy level diagram and absorption spectrum pair for  $\text{He}^+$ ?



D.)



E.)



For questions 4 and 5, refer to the energy level diagram for the hydrogen atom shown on page 1.

- 4.) Can hydrogen atoms in the ground (lowest energy) state absorb a photon of energy  $1.75 \times 10^{-18}$  J?
- A.) Yes, because hydrogen atoms can absorb any energy in the range from zero to  $2.18 \times 10^{-18}$  J.
  - B.) Yes, because the electron is promoted to an energy between the second and third energy levels. As the electron drops to the second level, the hydrogen atoms release the excess energy.
  - C.) Yes, because this represents the energy difference between the ground state and second energy level.
  - D.) No, because this much energy would cause the atom to be ionized.
  - E.) No, because this energy does not correspond to the energy difference between the ground state and any other energy level.
- 5.) Can hydrogen atoms in the ground (lowest energy) state absorb a photon of energy  $3.00 \times 10^{-18}$  J?
- A.) Yes, because this energy corresponds to the energy difference between the ground state and a quantized energy level with energy greater than zero.
  - B.) Yes, the energy of the light is greater than what is required to ionize the atom. The excess energy would be converted into kinetic energy.
  - C.) No, it is impossible for atoms to absorb light whose energy is greater than their ionization energy.
  - D.) No, atoms can only absorb light whose energy corresponds to the difference between the ground state and another level.
  - E.) No, light of this energy would not be enough to excite the electron from the ground state to the first energy level.

6.) Which has a higher ionization energy than Ne?

- A.) Ar      B.) F      C.)  $F^-$       D.)  $Na^+$       E.) Na

7.) How many unpaired electrons exist in the ground state electronic configuration  $[Ar]4s^23d^8$ ?

- A.) 0      B.) 1      C.) 2      D.) 3      E.) 4


8.) Which orbital has the quantum numbers  $n = 3$  and  $l = 1$ ?

- A.) 2s      B.) 2p      C.) 3s      D.) 3p      E.) 4d

9.) Identify X from the configuration  $X^+ (1s^22s^22p^63s^2)$ .

- A.) Na      B.) Mg      C.) Al      D.) Si      E.) P

10.) For one mole of an ideal gas at constant volume,  $X = ?$



- A.) molar mass      B.)  $v_{\text{rms}}$       C.)  $R$       D.)  $V$       E.)  $P$

11.) Absorption of what color light will induce the  $4 \rightarrow 6$  transition in  $\text{He}^+$ ?

- A.) Infrared (IR)      B.) Red      C.) Green  
D.) Blue      E.) Ultraviolet (UV)


**Part 2: Short Answer Problems (85 pts total)**

Instructions: Enter answers in the boxes provided. Show your work. Where requested write explanations in fifteen words or less.

**(40 pts)**

1.) Each figure in parts a-d contains at least two errors. In the space provided, specify two of the errors and provide a brief explanation of each one.


a.)



Error 1:

Error 2: .


b.)



Error 1:

Error 2:


c.)



Error 1:

Error 2:

d.)



Error 1:

Error 2:


**(45 pts)****2.)**

1) Ozone ( $O_3$ ) gas is placed in a 1.0 L glass vessel at a pressure of 2.0 atm and a temperature of 300 K. Assume ideal behavior.

a) What are the number of moles and the mass of ozone present?

Moles Ozone:
--------------

b) Shining ultraviolet light on  $O_3$  induces the following reaction:



If half of the ozone present reacts, what is the final mole fraction of each gas in the vessel?

Mole Fraction $O_2$ :
-----------------------

Mole Fraction $O_3$ :
-----------------------



c.) Calculate the partial pressure of each gas and the total pressure.

Pressure O <sub>3</sub> :
Total Pressure:

d) The graph below depicts the speed distribution of O<sub>3</sub> molecules at 300K. Using the same axes, sketch the distribution of O<sub>3</sub> at 1200K.