MCB135E, 1st midterm October 2, 1998

Name: SID:

I. (45 points) Multiple choice questions. Only one answer is correct, 3 points each correct answer.

- 1. Growth and development are processes that:
 - A. continue throughout the lifespan
 - B. follow a precise, predictable timetable
 - C. can be influenced by genetic and environmental factors
 - D. can affect function during maturity and length of lifespan
 - E. all of the above
- 2. Ovulation depends on the following events, EXCEPT:
 - A. occurs at approximately the middle of the menstrual cycle
 - B. depends on high circulating levels of estrogens
 - C. depends on high circulating levels of progesterone
 - D. depends on LH surge
 - E. depends on high levels of GnRH
- 3. Fertilization occurs:
 - A. in the oviduct, close to the ovary
 - B. in the oviduct, close to the uterus
 - C. ovum can be fertilized as long as 48 hours after ovulation
 - D. ovum looses its capacity to be fertilized 10 hours after ovulation
 - E. A & C
- 4. Sperm and ovum resemble in many characteristics. Some of the major similarities are EXCEPT:
 - A. they have an haploid number of chromosomes
 - B. they derive from primitive cells by the processes of mitosis/meiosis
 - C. after puberty, sperm and ovum production continues throughout life
 - D. maturation of sperm and ovum depends on sex hormones
 - E. maturation of sperm and ovum depends on gonadotropins
- 5. At fertilization, the major functions of the sperm include all of the following, EXCEPT:
 - A. it reaches and penetrates the ovum
 - B. activates the ovum to nuclear and cytoplasmic division to achieve the haploid number of maternal chromosomes
 - C. provides food reserves for growth and development of zygote
 - D. contributes the paternal complement of chromosomes
 - E. determines the sex of the embryo

Name: SID:

- 6. The Sertoli cells,
 - A. are found along the wall of the seminiferous tubules
 - B. secrete chemical messengers for sperm production/maturation
 - C. phagocytize defective sperm
 - D. are stimulated by androgens and FSH
 - E. all of the above
- 7. The Turner's syndrome or ovarian dysgenesis is characterized by the following signs, EXCEPT:
 - A. the sex chromosomes are XO
 - B. the ovary is underdeveloped and hypofunctioning
 - C. high stature
 - D. cardiac abnormalities
 - E. some deficits of the nervous system
- 8. Until the 6th embryonal week, the primitive gonad:
 - A. is undifferentiated
 - B. is differentiated into a male type
 - C. is differentiated into a female type
 - D. is able to produce sex hormones
 - E. none of the above
- 9. The SRY gene or group of genes on the Y chromosome is responsible for inducing a cascade of differentiating events leading to:
 - A. the differentiation of the Leydig cells of the testis
 - B. the production of testosterone
 - C. the production of the Mullerian inhibiting substance by the Sertoli cells
 - D. the masculinization of the male secondary sex organs
 - E. all of the above
- 10. Of the major CNS structures derived from the embryonal vesicles, those derived from the telencephalon include all of the following EXCEPT:
 - A. cerebral cortex
 - B. limbic system
 - C. hypothalamus
 - D. corpus striatum
 - E. retina
- 11. In the human placenta, the following statements are correct, EXCEPT:
 - A. maternal and fetal blood mix in the maternal sinuses
 - B. maternal and fetal blood do not enter in direct contact but are separated by the wall of the floating villi
 - C. exchange of gases occurs by diffusion from maternal to fetal blood
 - D. delivery of nutrients occurs from mother to fetus
 - E. removal of waste materials occurs from fetus to mother

Name: SID:

12. The placenta reaches its peak (most optimal) function at midgestation, from then on the placenta:

A. undergoes aging changes

B. is less able to protect the fetus from environmental toxins

C. may be responsible for insufficient fetal growth (small-for-date newborns)

D. may be responsible for a decline in hormonal production

E. all of the above

13. The following statements are correct EXCEPT:

A. in the CNS, myelin formation depends on oligodendrocytes

B. oligodendrocytes are glial cells

C. myelin is essential for rapid conduction of nerve impulses

D. myelin formation is exclusivly a prenatal process

E. myelin formation can continue until the 3rd decade of life

14. Certain features characterize the different age periods. Which of the following periods is INCORRECTLY matched to its characteristics?

A. embryonic period: first trimester of prenatal life with rapid differentiation and establishement of organ systems

B. early fetal period: second trimester of prenatal life with accelerated growth and elaboration of structures

C. late fetal period: third trimester of prenatal life with rapid increase in body mass and cessation of placental function

D. neonatal period, newborn: birth to end of second postnatal week

E. pubertal period of adolescence: secondary sex organs maturation

15. In the fetus, the major source of eenergy derives from:

A. sugar and carbohydrates

B. proteins

C. Lipids

D. none of the above

E. all of the above

Il (14 points). True (A) False (B) 2 points each correct answer

16. Androgens (such as testosterone) 'activate'male behavior prenatally and 'organize' CNS structures postnatally

17. Lordosis, a special posture in female rats designed to favor copulation, is dependent on the presence of estrogens (estradiol) during neonatal days 1-4

Name:
SID:
18. In neonates, total daily REM sleep is longer than in adults
19. Growth hormone (GH) from the pituitary is present in the fetus and is necessary for fetal growth
20. Newborns see colors, and by 3-4 months postnatally, infants can distinguish colors as well as adults.
21. The newborn is significantly more susceptible to hypoxia than the adult due to its reliance on glycolytic respiration
22. The 3 germ layers of the gastrula (ectoderm, mesoderm and endoderm) will eventually differentiate into all the embryo's tissues and organs.
III. (21 points)Describe the 3 fetal structures that are unique to the fetal circulation:
23. (6 points) Name them (a.) DUCTUS VENOSUS (b.) FORAMEN OVALE (c.) FORAMEN OVALE 24. (6 points) Indicate their connections and the blood oxygen/nutrients content (high, medium or low?) that they carry.
(a.) Carries blood with High oxygen/nutrient content from the UMBELICAL VEIN to the TYC.
(b.) Carries blood with to the to the Cack
(c.) Carries blood with MED) exygen/nutrient content from the to the AORTA 25. (3 points) Explain in a few sentences their physiological significance
2 P+5 growth. The second of the fetal circulation and fetal
SHUNTING OF OXYGENATED BIOOD OF Brain,
SHUNTING OF DXYGENATED Blood To Organs that have PRIORITY: LIVER, & Brain, head & NECK them the lower Body
AMY OF HIS 3 STRUCTURE JOES Its

Name: SID:

IV (20 points, 2 points each correct answer) Match the hormone/endocrine conditions on the left with the appropriate functions/characteristics on the right. Use each answer once.

- B_26. testosterone
- A 27. human chorionic gonadotropin
- 28. human chorionic somatomammotropin C. Is necessary for
- E__29. Mullerian inhibitor substance
- F__30. prenatal thyroid insufficiency
- __31. 5-alpha reductase
- (<u>___32</u>. dihydrotestosterone
- 33. progesterone
 - I 34. estradiol
- ## 35. aldosterone

- A. Maintains corpus luteum and its hormonal secretions
- B. Is secreted by the fetal testis by the 6-8 embryonal weeks
- C. Is necessary for conversion to dihydrotestosterone
- D. Regulates maternal metabolism
- E. Induces regression of Mullerian ducts
- F. Cretinism
- G. Is necessary for differentation of the male internal genitalia
- H. Is one of the hormones of the fetal adrenal cortex with low levels until late pregnancy
- I. Is produced by the thecal cells from the ovarian follicle
- J. Is necessary for preparing the uterus for implantation of the blastocyst.