

J. Strain

Math 128A final exam 24 May 2003 ①

1. (a) Find the Newton form of the quadratic polynomial interpolating  $f(x) = \ln x$  at  $x = 1, e$  and  $e^2$ .

(b) Show that the error is no larger than 512 for  $1 \leq x \leq e^2$ .

(c) Evaluate the actual error at  $x = 5$ .

2. Let

$$A = \begin{bmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}.$$

(a) Find a lower triangular matrix L and an upper triangular matrix U such that  $PA = LU$  for some permutation matrix P. Explain your procedure.

(b) Is A diagonally dominant? Positive definite? Does either answer affect the procedure and result of (a)?

(c) Is A invertible? Why or why not?

3.(a) For the initial value problem <sup>(2)</sup>

$$y' = f(t, y), \quad y(0) = y_0$$

define the local truncation error  $T_n$  of a numerical method

$$u_{n+1} = u_n + h \Phi(t_n, h; u_n, \dots, u_{n-k}; f).$$

(b) Show that for the trapezoidal rule

$$u_{n+1} = u_n + h \left[ \frac{1}{2} f(t_{n+1}, u_{n+1}) + \frac{1}{2} f(t_n, u_n) \right]$$

the local truncation error is  $O(h^2)$ .

(c) How can you use the trapezoidal rule (with various step sizes  $h$ ) to construct a numerical method with local truncation error  $O(h^4)$ ?

4. Determine the limit and speed of convergence for the iteration

$$x_{n+1} = 2x_n - \pi x_n^2, \quad x_0 = \sqrt{3}.$$

Use analysis (not a calculator).

5. Suppose  $u \in \mathbb{R}^n$  is a unit vector  
 $(\|u\|_2 = 1)$ .

(3)

(a) Explain the geometric meaning of the matrices

$$Q = uu^T, \quad P = I - uu^T, \quad R = I - 2uu^T.$$

(b) Find a  $3 \times 3$  orthogonal matrix  $Q$  such that

$$Q \begin{bmatrix} 3 \\ 0 \\ 4 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 0 \end{bmatrix}.$$


6.(a) Find the weights  $w_i$  of a quadrature formula

$$\int_0^1 f(x)dx \approx w_0 f(0) + w_1 f(1) + w_2 f'(1)$$


which has as high precision as possible.

(b) Without any additional calculation, find an equally accurate formula of the form

$$\int_0^1 f(x)dx \approx w_0^* f(0) + w_1^* f(1) + w_2^* f'(0).$$



(c) Consider compounding the rules of (a) and (b) by applying them alternately on intervals of length  $h = \frac{1}{2n}$  from 0 to 1:



Write out the resulting rule in the form

$$\int_0^1 f(x) dx = a_0 f(0) + a_1 f(h) + a_2 f(2h) + \dots + a_{2n} f(2nh) + b_0 f'(0) + b_1 f'(h) + \dots + b_{2n} f'(2nh).$$

(d) Find the largest integer  $p$  such that the error of the rule in (c) is

$$O(h^p),$$

and determine by dimensional analysis the order  $q$  of differentiation that will make the error  $O(f^{(q)} h^p)$ .