


**SECTION 1: EQUILIBRIUM**




For questions 1 – 11 consider the following three reactions at 298 K.



I)  $\frac{[\text{S}]^2 [\text{O}_2]^3}{[\text{SO}_3]^2} = 0.225$

$\frac{[\text{SO}_2]^2}{[\text{S}] [\text{O}_2]} = 225 \cdot 5.062 \cdot 10^4$

$K = \frac{[\text{SO}_2]^2 [\text{O}_2]}{[\text{SO}_3]^2}$



1.) What is the equilibrium constant for a mixture of  $\text{O}_2$ ,  $\text{SO}_3$  and  $\text{SO}_2$  gas (rxn. III)?

- A) 75.5
- B) 112
- C)  $1.1 \times 10^4$
- D)  $2.5 \times 10^5$
- E) 0.775

2.) What is the value of the equilibrium constant for rxn I if at equilibrium the flask contains 0.236 atm  $\text{SO}_3$ , 0.500 atm  $\text{O}_2$ , and 0.01 g Sulfur after a temperature change.

- A) 0.0909
- B) 11.0
- C)  $1.63 \times 10^{-5}$
- D)  $6.25 \times 10^{-2}$
- E) 2.24

3.) What change has occurred if the value of K for rxn I is found to be 0.552?

- A) An increase in temperature.
- B) A decrease in temperature.
- C) An increase in pressure.
- D) An increase in volume.
- E) cannot be determined.

4.) Which is a suitable expression for the reaction quotient for the formation of  $\text{SO}_2$  from the elements?


- A)  $P(\text{O}_2) / P(\text{SO}_2)$
- B)  $P(\text{O}_2) P^2(\text{SO}_2) / P(\text{S})$
- C)  $P(\text{O}_2) / P^2(\text{SO}_2) P(\text{S})$
- D)  $P(\text{SO}_2) / P(\text{O}_2)$
- E) Nothing can be said with the information given.

$\frac{P_{\text{SO}_2}}{P_{\text{O}_2}}$

5.) What is  $\Delta H^\circ$  for reaction III?

- A) 333 kJ
- B) 251 kJ
- C)  $1.7 \times 10^3 \text{ kJ}$
- D) 5 kJ
- E) 76 kJ

6.) Which is the best arrangement of the relative enthalpies of formation of compounds O<sub>2</sub>, SO<sub>3</sub>, and SO<sub>2</sub>?



$$\begin{aligned} \text{SO}_3 &= -395.5 \\ \text{SO}_2 &= -270 \\ \text{O}_2 &= 0 \end{aligned}$$

7.) What is the best prediction of  $\Delta S^\circ$  for reaction I at 298K?

- A)  $\Delta S^\circ > 0$
- B)  $\Delta S^\circ = 0$
- C)  $\Delta S^\circ < 0$
- D)  $\Delta S^\circ \leq 0$
- E)  $\Delta S^\circ \geq 0$


$$\begin{aligned} \Delta H &\oplus \\ \Delta S &\oplus \end{aligned}$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$$

8.) What can you say about reaction I at 298 K?

- A) It is exothermic.
- B) It is spontaneous.
- C) It is not spontaneous.
- D) It is at equilibrium.
- E) It releases heat.

9.) The correct plot for  $\ln K$  vs  $1/T$  for reaction I would pass through which pair of points (fill in both points on scantron sheet)?



$$\ln K = -\frac{\Delta H^\circ}{R} \left( \frac{1}{T} \right) + \frac{\Delta S^\circ}{R}$$

10.) From which of the following starting conditions would it be impossible for equilibrium to be achieved for reaction II?


- A) Pure SO<sub>2</sub> (g).
- B) A mixture of SO<sub>2</sub> (g), O<sub>2</sub> (g), and S (s).
- C) A mixture of SO<sub>2</sub> (g) and O<sub>2</sub> (g).
- D) Pure O<sub>2</sub> (g) and S (s).
- E) Equilibrium can be achieved from any of these starting conditions.

11.) Which occurs when adding S (s) to the equilibrium described by reactions I, II and III?

- A) A decrease in the pressure of SO<sub>3</sub>(g).
- B) A decrease the pressure of SO<sub>2</sub>(g).
- C) An increase in the value of the equilibrium constant.
- D) An increase in the total pressure of the system.
- E) No change in the equilibrium.

Continue with the next question:


12.) For the reaction



having  $\Delta G^\circ = -2400 \text{ kJ}$  at  $25^\circ\text{C}$ , the equilibrium mixture \_\_\_\_\_.

- A) will consist almost exclusively of A and D.
- B) will consist almost exclusively of A and Z.
- C) will consist almost exclusively of X and Z.
- D) will consist of significant amounts of A, D, X, and Z.
- E) has a composition predictable only if one knows T and  $\Delta H^\circ$  and  $\Delta S^\circ$ .

13.) The equilibrium constant for the reaction below at  $25^\circ\text{C}$  is  $4.8 \times 10^{-6}$ . Calculate the equilibrium concentration (mol/L) of Cl<sub>2</sub> (g) if the initial concentration of ICl (g) is 1.33 mol/L. There is no I<sub>2</sub> or Cl<sub>2</sub> initially present.



- A)  $2.9 \times 10^{-3}$
- B)  $5.8 \times 10^{-3}$
- C)  $3.2 \times 10^{-6}$
- D)  $6.4 \times 10^{-6}$
- E) 343

$$4.8 \times 10^{-6} = \frac{[\text{I}_2][\text{Cl}_2]}{[\text{ICl}]^2} = \frac{x^2}{(1.33-x)^2}$$

$$2.19 \times 10^{-3} = \frac{x}{1.33-x}$$


$$x = 2.91 \times 10^{-3}$$

14.) Which of the following equilibria, will shift to the left in response to a decrease in volume?

- A)  $\text{H}_2(g) + \text{Cl}_2(g) \leftrightarrow 2 \text{HCl}(g)$
- B)  $2 \text{SO}_3(g) \leftrightarrow 2 \text{SO}_2(g) + \text{O}_2(g)$
- C)  $\text{N}_2(g) + 3 \text{H}_2(g) \leftrightarrow 2 \text{NH}_3(g)$
- D)  $4 \text{Fe}(s) + 3 \text{O}_2(g) \leftrightarrow 2 \text{Fe}_2\text{O}_3(s)$
- E)  $2\text{HI}(g) \leftrightarrow \text{H}_2(g) + \text{I}_2(g)$

fewer mols

15.) Consider the following reaction at equilibrium:



How can the yield of  $\text{CO}(\text{g})$  be maximized?

- A) at high temperature and high pressure
  - B) at high temperature and low pressure
  - C) at low temperature and low pressure
  - D) at low temperature and high pressure
  - E) in the presence of solid carbon
- 16.) Which is true for every reaction if the temperature is raised?
- A) Chemical reactions favor products.
  - B) Chemical reactions favor reactants.
  - C) No change is observed.
  - D) Equilibrium constants increase.
  - E) None of these.

For the following three questions, consider the equilibrium  $\text{PbSO}_4(\text{s}) \leftrightarrow \text{Pb}^{2+}(\text{aq}) + \text{SO}_4^{2-}(\text{aq})$  which has a  $K_{\text{sp}} = 1.6 \times 10^{-8}$  at 298 K

17.) What is the concentration of lead ions in water (M) when solid  $\text{PbSO}_4$  is present?

- A)  $1.6 \times 10^{-10}$
- B)  $1.3 \times 10^{-4}$
- C) 1.0
- D)  $1.1 \times 10^4$
- E) 22.5

18.) What is the concentration of lead ions (M) in 0.01 M  $\text{Na}_2\text{SO}_4$  ( $K_{\text{sp}} \sim 10^{-8}$ ) when solid  $\text{PbSO}_4$  is present?

- A)  $1.6 \times 10^{-6}$
- B)  $1.3 \times 10^{-4}$
- C) 1.0
- D)  $1.1 \times 10^4$
- E) 22.5

$$(0.01 + x)x \approx 1.6 \times 10^{-8}$$

19.) What is  $\Delta G^\circ$  for the dissolution of lead sulfate at 298 K (kJ/mol)?

- A) -44
- B) -13
- C) 5.9
- D)  $1.1 \times 10^4$
- E)  $2.3 \times 10^{-3}$


$$\ln(K) = -RT$$

$$\Delta G^\circ = -RT \ln K = \frac{-8.3145 \times 10^{-3} \text{ kJ} (298 \text{ K}) \ln(1.6 \times 10^{-8})}{\text{mol} \cdot \text{K}}$$

Continue with the next question:

**SECTION 2: PHASES OF MATTER**

For the following questions consider the phase diagram for water below.



20.) At which point are gas, liquid and solid all in equilibrium?

- A) (T2, P2)
- B) (T2, P1)
- C) (T1, P1)
- D) (T3, P1)
- E) (T3, P3)

21.) Arrow I corresponds to:

- A) Constant pressure
- B) Equilibrium
- C) Sublimation
- D) Condensation
- E) Melting

22.) Along the curve containing the points (T2, P2) and (T3, P3):

- A) Solid, liquid and gas are all in equilibrium.
- B) Liquid and gas are in equilibrium.
- C) The vapor pressure is constant.
- D) The gas cannot be condensed at any pressure.
- E) Only the solid phase is observed.

23.) At the point (T2, P3) the substance is:

- A) In equilibrium between liquid and gas.
- B) A liquid.
- C) A gas.
- D) A supercritical fluid.
- E) A solid.

24.) Which is true at temperatures above  $T_3$ ?

- A) Solid, liquid and gas are all in equilibrium.
- B) Liquid and gas are in equilibrium.
- C) The vapor pressure is constant.
- D) The gas cannot be condensed at any pressure.
- E) Only the solid phase is observed.

25.) Which intermolecular force predominates in the condensation of water?

- A) H-bonding
- B) Van der Waals
- C) London
- D) Ion-Ion
- E) Dipole-Ion

Continue with the next question:

### SECTION 3: THERMODYNAMICS

26.) Which one of the following is always positive when a spontaneous process occurs?

- A)  $\Delta S_{\text{system}}$
- B)  $\Delta S_{\text{surroundings}}$
- C)  $\Delta S_{\text{universe}}$
- D)  $\Delta H_{\text{universe}}$
- E)  $\Delta H_{\text{surroundings}}$

$$\Delta G^\circ = \Delta H - T\Delta S$$

27.) Which is true of the entropy of the universe?

- A) conserved.
- B) continually decreasing.
- C) continually increasing.
- D) equal to zero.
- E) equal to the energy, E.

28.) Which is a state function (mark all that apply)?

- A) flame heating.
- B) enthalpy.
- C) entropy.
- D) electrical work.
- E) none of these.

29.) What is the change in the internal energy (in J) of a system that releases 1000 J of heat and does 225 J of work on the surroundings?

- A) -10,155
- B) 1225
- C) -775
- D) 775
- E) 1225

$$\begin{aligned}\Delta U &= q + w \\ &= -1000 - 225\end{aligned}$$

30.) What do you expect the temperature change to be for the rapid, adiabatic compression of a gas from 1.0 atm to 3.0 atm?

- A) -10K
- B) 100K
- C) 0.001K
- D) -100K
- E) -0.001K

$$W = -P\Delta V \quad q = 0 \quad W = \Delta T$$

$$m = nRT$$

$$\frac{P_1}{V_1} = \frac{P_2}{V_2} \quad \frac{V_2}{V_1} = \frac{P_1}{P_2} = \frac{1}{3}$$

31.) A bar of hot metal is placed in water in an insulated container and the two are allowed to reach thermal equilibrium. When 1.0 kg of metal at 100°C is placed in 2.0 kg of water, the temperature water bath raises from 20°C to 25°C. What is the specific heat capacity of the metal (J/g K)?

- A) 0.5
- B) 1.5
- C) 0.22
- D) 25
- E) .025

$$m_{\text{metal}} c \Delta T = m_{\text{H}_2\text{O}} c \Delta T$$

$$c = \frac{m_{\text{H}_2\text{O}} c \Delta T}{m_{\text{metal}} \Delta T} = \frac{2.0 \text{ kg} \times 4.184 \text{ J/g K} (5 \text{ K})}{1.0 \text{ kg} (75)}$$

32.) Which is the best estimate for the boiling point of benzene (°C) given that  $\Delta H^\circ$  of vaporization is 31 kJ/mol and  $\Delta S^\circ$  of vaporization is 90 J/mol K?

- A) 25
- B) 45
- C) 65
- D) 15
- E) 5

344 K


$$\Delta S^\circ = \frac{\Delta H^\circ}{T}$$

$$T = \frac{\Delta H^\circ}{\Delta S^\circ} = \frac{31 \text{ kJ/mol}}{90 \text{ J/mol K} \times \frac{1 \text{ kJ}}{1000 \text{ J}}} = 344 \text{ K}$$

33.) Which is the first step in a realistic experiment to determine the entropy change for a chemical reaction?

- A) Measuring  $\Delta H^\circ$ .
- B) Counting the microstates.
- C) Counting the change in microstates.
- D) Measuring how the K varies with temperature.
- E) The entropy change cannot be measured for chemical reactions.

34.) The value of  $\Delta H^\circ$  for the following reaction is -3351 kJ.



What is  $\Delta H^\circ$  for the formation of 75.0 g of  $\text{Al}_2\text{O}_3(s)$  (kJ)?

- A)  $-2.51 \times 10^7$
- B)  $-1.26 \times 10^5$
- C) -2460
- D) -1230
- E) +3351

$$75.0 \text{ g} \times \frac{1 \text{ mol Al}_2\text{O}_3}{101.96 \text{ g}} \times \frac{-3351 \text{ kJ}}{2 \text{ mol Al}_2\text{O}_3} = -1230 \text{ kJ}$$

30.) What do you expect the temperature change to be for the rapid, adiabatic compression of a gas from 1.0 atm to 3.0 atm?

- A) -10K
- B) 100K
- C) 0.001K
- D) -100K
- E) -0.001K

$$W = -P\Delta V \quad q = 0 \quad W = \Delta T$$

$$P_1 V_1 = P_2 V_2 \quad m = nRT$$

$$\frac{P_1}{V_1} = \frac{P_2}{V_2} \quad \frac{V_2}{V_1} = \frac{P_1}{P_2} = \frac{1}{3}$$

31.) A bar of hot metal is placed in water in an insulated container and the two are allowed to reach thermal equilibrium. When 1.0 kg of metal at 100°C is placed in 2.0 kg of water, the temperature water bath raises from 20°C to 25°C. What is the specific heat capacity of the metal (J/g K)?

- A) 0.5
- B) 1.5
- C) 0.22
- D) 25
- E) .025

$$m_{\text{metal}} c \Delta T = m_{\text{H}_2\text{O}} c \Delta T$$

$$c = \frac{m_{\text{H}_2\text{O}} c \Delta T}{m_{\text{metal}} \Delta T} = \frac{2.0 \text{ kg} \times 4.184 \text{ J/g K} (5 \text{ K})}{1.0 \text{ kg} (75)}$$

32.) Which is the best estimate for the boiling point of benzene (°C) given that  $\Delta H^\circ$  of vaporization is 31 kJ/mol and  $\Delta S^\circ$  of vaporization is 90 J/mol K?

- A) 25
- B) 45
- C) 65
- D) 15
- E) 5

344 K


$$\Delta S^\circ = \frac{\Delta H^\circ}{T}$$

$$T = \frac{\Delta H^\circ}{\Delta S^\circ} = \frac{31 \text{ kJ/mol}}{90 \text{ J/mol K} \times \frac{1 \text{ kJ}}{1000 \text{ J}}} = 344 \text{ K}$$

33.) Which is the first step in a realistic experiment to determine the entropy change for a chemical reaction?

- A) Measuring  $\Delta H^\circ$ .
- B) Counting the microstates.
- C) Counting the change in microstates.
- D) Measuring how the K varies with temperature.
- E) The entropy change cannot be measured for chemical reactions.

34.) The value of  $\Delta H^\circ$  for the following reaction is -3351 kJ.



What is  $\Delta H^\circ$  for the formation of 75.0 g of  $\text{Al}_2\text{O}_3(s)$  (kJ)?

- A)  $-2.51 \times 10^7$
- B)  $-1.26 \times 10^5$
- C) -2460
- D) -1230
- E) +3351

$$75.0 \text{ g} \times \frac{1 \text{ mol Al}_2\text{O}_3}{101.96 \text{ g}} \times \frac{-3351 \text{ kJ}}{2 \text{ mol Al}_2\text{O}_3} = -1230 \text{ kJ}$$




35.) Which of the following has a non-zero  $\Delta H_f^\circ$ ?

- A)  $O_2(l)$
- B) C(graphite)
- C)  $N_2(g)$
- D)  $F_2(g)$
- E)  $Cl_2(g)$

36.) Which one of the following processes is endothermic?

- A)  $2H_2(g) + O_2(g) \rightarrow 2H_2O(g)$
- B)  $H_2O(g) \rightarrow H_2O(l)$
- C)  $CH_4(g) + 2O_2(g) \rightarrow CO_2(g) + 2H_2O(g)$
- D)  $H_2O(s) \rightarrow H_2O(l)$
- E)  $2Al(s) + Fe_2O_3(s) \rightarrow Al_2O_3(s) + 2Fe(l)$

37.) Which is true for the following reaction under standard conditions?




- A) spontaneous at all temperatures
- B) spontaneous only at high temperature
- C) spontaneous only at low temperature
- D) not spontaneous at all temperatures
- E) cannot be determined

$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$   
 $= 137 - 120T$   
 spon @ high temp

38.) Given the following

Substance	$\Delta H_f^\circ$ (kJ/mol)
$SO_2(g)$	-297
$SO_3(g)$	-396
$SO_2Cl_2(g)$	-364
$H_2SO_4(l)$	-814
$H_2O(l)$	-286

Calculate the amount of heat (in kJ) evolved when 11.25 g of  $SO_2$  reacts according to the equation:




- A) 100.5
- B)  $8.550 \times 10^5$
- C) 47.5
- D) 11.25
- E) Insufficient data are given.


$11.25g \times \frac{1 \text{ mol } SO_2}{64.07g} \times \frac{-67}{1 \text{ mol } SO_2}$   
 $\Delta H^\circ = -67 \text{ kJ/mol}$

39.) We have seen many times in lecture that heat is given off in the combustion of hydrogen gas. Which of the following is responsible for the heat?

- A) Breaking H-H and O-O bonds.
- B) Breaking O-H bonds.
- C) Forming H-H bonds and O-O bonds.
- D) Forming O-H bonds.
- E) Condensation of the water formed.



40.) What can you say about the reaction if the ratio of the C=C double bond strength to the C-C single bond strength is less than two?



break 2 double bonds < 4 single bonds  
⊕ E

du

- A) The reaction is exothermic.
- B) The reaction is endothermic.
- C) The reaction is spontaneous.
- D) The enthalpy change is about zero.

1 double bond weaker than 2 single

bond E broken - bonds formed

broken < f.

⊖

exo.