




Chemistry 1A Fall 1999



Midterm Exam III, version A

November 9, 1999

(Closed book, 90 minutes, 145 points)

Name: _____

Section Number: _____

SID: _____

T.A. Name: _____

Identification Sticker

Exam information, exam directions, and useful hints to maximize your score:

- ▶ Write your name on all 6 pages.
- ▶ There are two parts to this exam: 1) multiple choice and 2) short answer problems.
- ▶ **For the multiple choice problems, fill in the Scantron™ form AND circle the answer on your exam.**
- ▶ Answer the questions you know how to do first, then work on the questions you skipped.
- ▶ Show all work for which you want credit and do not forget to include units!
- ▶ You may use the back side of the exam pages to show your work and/or for scratch paper.

Potentially useful information:

Unit Prefixes

milli, m ($\times 10^{-3}$)	micro, μ ($\times 10^{-6}$)	nano, n ($\times 10^{-9}$)
kilo, k ($\times 10^3$)	mega, M ($\times 10^6$)	giga, G ($\times 10^9$)

$$\Delta G^\circ = \sum \Delta G_f^\circ(\text{products}) - \sum \Delta G_f^\circ(\text{reactants})$$

$$\Delta H^\circ = \sum \Delta H_f^\circ(\text{products}) - \sum \Delta H_f^\circ(\text{reactants})$$

$$\Delta S^\circ = \sum S^\circ(\text{products}) - \sum S^\circ(\text{reactants})$$

$$c_p(\text{H}_2\text{O}(l)) = 75.3 \frac{\text{J}}{\text{mol} \cdot \text{K}}$$

$$q_v = nc_v \Delta T \quad q_p = nc_p \Delta T$$

$$\Delta G = \Delta H - T\Delta S \quad S = k_B \ln W$$

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ = -RT \ln K$$

$$\Delta E = q + w \quad PV = nRT$$

$$1 \text{ cal} = 4.184 \text{ J} \quad 1000 \text{ cal} = 1 \text{ Cal}$$

(Do not write in this box, it's for official use only)

Thermodynamic Properties		
Substance	ΔH_f° (kJ/mol)	S° (J/mol·K)
Cl ₂ (g)	0.0	223.0
CO ₂ (g)	-393.5	213.6
H ₂ O (l)	-285.8	69.91
H ₂ SO ₄ (l)	-814.0	156.9
Fe ₂ O ₃ (s)	-824.2	87.4
FeS ₂ (s)	-178.2	52.93
O ₂ (g)	0.0	205.03

Problems	Points
multiple choice	/ 44
Part 2, # 1	/ 30
Part 2, # 2	/ 18
Part 2, # 3	/ 25
Part 2, # 4	/ 28
all	/ 145

Page 2 of 6


Name: _____

Part 1: Multiple Choice.**(4 pts each, 44 pts total)**

Instructions: Bubble in the correct answer on your Scantron™ form AND circle the answer on your exam. Each question has one correct answer.

1.) The answer to question 1 is A. Bubble in A on your Scantron™ form.

2.) Consider the sublimation of dry ice:



If K_1 is the equilibrium constant at 300 K, and K_2 is the equilibrium constant at 400 K, which of the following inequalities must be true?

- A.) $K_1 = K_2$ B.) $K_1 = K_2^{-1}$ C.) $K_1 K_2 = 0$ D.) $K_1 > K_2$ E.) $K_1 < K_2$


3.) For the vaporization of methanol



$\Delta H^\circ = 38.0 \text{ kJ mol}^{-1}$ and $\Delta S^\circ = 112.9 \text{ J K}^{-1} \text{ mol}^{-1}$. What is the boiling point of methanol at sea level? Assume ΔH° and ΔS° are independent of T.

- A.) 64 K B.) 237 K C.) 273 K D.) 337 K E.) 373 K

4.) Consider the reaction:



at equilibrium. What would be the reaction quotient immediately following the reduction of volume by two at constant temperature before any reaction occurs?

- A.) $Q = \frac{1}{4}K$ B.) $Q = \frac{1}{2}K$ C.) $Q = K$ D.) $Q = 2K$ E.) $Q = 4K$

5.) One mole of an ideal gas expands *isothermally* against a constant pressure of 1 atmosphere. Which of the following inequalities is true?


- A.) $\Delta P > 0$ B.) $q > 0$ C.) $\Delta S < 0$ D.) $\Delta V < 0$ E.) $\Delta T < 0$

Page 3 of 6

Name: _____

- 6.) One mole of an ideal gas expands *adiabatically* against a constant pressure of 1 atmosphere. Which of the following inequalities is true?
- A.) $\Delta P > 0$ B.) $q > 0$ C.) $\Delta S < 0$ D.) $\Delta V < 0$ E.) $\Delta T < 0$
- 7.) At what temperature does $K = 1$, $\Delta G^\circ = 0$ for the reaction $\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_2\text{O}(\text{g})$?
- A.) -273°C B.) 0°C C.) 100°C D.) 273°C E.) 373°C
- 8.) How many different ways can you distribute six distinguishable stones between two boxes with five in the first box and one in the second box?
- A.) 1 B.) 3 C.) 6 D.) 9 E.) 15
- 9.) The caloric content of 10 little cookies can heat up 10 kg of water by 10°C . What would be the change in temperature if 1 little cookie was used to heat up 1 kg of water?
- A.) 0.1°C B.) 1.0°C C.) 10°C D.) 100°C E.) 1000°C

For each of the problems 10-12, select the graph that best describes the behavior listed.



- 10.) $P_{\text{N}_2\text{O}_4}$ as a function of $(P_{\text{NO}_2})^2$ for $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$, at constant T .
- 11.) $\ln(K)$ as a function of $\frac{1}{T}$ for the combustion of liquid methanol (CH_3OH).
- 12.) ΔG° as a function of T for the vaporization of water, $\text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{H}_2\text{O}(\text{g})$.

Page 4 of 6

Name: _____

Part 2: Short Answer Problems (101 pts total)

Instructions: Enter answers in the boxes provided. Show your work. Explain your answer when requested in 15 words or less.

(30 pts)

1.) The reaction

is endothermic with $\Delta H = 4.6 \text{ kJ/mol}$. $\text{SO}_2\text{Cl}_2(\text{g})$ and $\text{SO}_2(\text{g})$ are placed in a bulb at a fixed temperature with partial pressures of 3.0 atm each.

a) Write the expression for reaction quotient (Q) and calculate its value before any reaction occurs.

Answers:

b) After equilibrium is reached in the bulb at the same fixed temperature, the partial pressure of Cl_2 (P_{Cl_2}) is found to be 1.0 atm. What are the partial pressures of SO_2Cl_2 and SO_2 ($P_{\text{SO}_2\text{Cl}_2}$ and P_{SO_2})?

Answers:

c) Calculate the value of the equilibrium constant for the reaction in part b).

Answer:

d) If some Cl_2 is added to the equilibrium mixture, will the pressure of SO_2 increase, decrease, or stay constant as the system approaches the new equilibrium state? Circle the answer and explain.

Decrease

Same

Increase


Explanation:

Page 5 of 6

Name: _____

(18 pts)

- 2.) Consider the reaction of silver chloride (AgCl) dissolving in water.



- a) Dissolving 14.3 g of AgCl (s) consumes 6.5 kJ of heat. What is the temperature change if 14.3 g of AgCl(s) totally dissolves in 1.00 L of water initially at 20 °C?


Answer:

- b) In actuality the equilibrium constant (K) for this reaction is very small (1.6×10^{-10} at 25.0 °C). How will this affect the temperature change predicted in part (a)? Explain.

Answer:

(25 pts)

- 3.) A study of the geology of the earth shows that rocks older than 2 billion years contain iron in the form of FeS₂. In rocks less than 2 billion years old, iron appears mostly as the oxide Fe₂O₃ (hematite).



- a) Calculate ΔH° for the above reaction.

Answer:

- b) Calculate ΔS° for the above reaction.

Answer:


- c) Over what temperature range is this reaction spontaneous? As always, show your calculations.

Answer:

Page 6 of 6

Name: _____

(28 pts)

4.) Consider the hydrogenation of formaldehyde ($\text{H}_2\text{C}=\text{O}$) to form methanol (CH_3OH).

Average Bond Energy (kJ/mol)			
H-H	436	C-O	360
H-C	413	C=O	743
H-O	463	C-C	348
O-O	146	C=C	612
O=O	497	C≡C	838

a) Estimate ΔH° for this reaction.

Answer:

b) The *formation* of which species, formaldehyde or methanol, is more exothermic (i.e. has the lower ΔH_f°)?

Answer:

c) The *combustion* of which species, formaldehyde or methanol, produces more heat per mole?

Answer: