

## Chemistry 3A – Spring 2002

## Final Exam

Professor Jean Fréchet

Your full signature\_\_\_\_\_

May 20, 2002

Print your full name\_\_\_\_\_

Your SID\_\_\_\_\_

*Please check the section number and name of your GSI/TA.*

\_\_\_\_\_ 161 Margot Paulick

\_\_\_\_\_ 171 Danielle Dube

\_\_\_\_\_ 181 Matt Pratt

\_\_\_\_\_ 191 Laurie Schenkel

\_\_\_\_\_ 111 Cathleen Yung

\_\_\_\_\_ 121 Priya Sonik

\_\_\_\_\_ 131 Paul Furuta

\_\_\_\_\_ 141 Zach Fresco

\_\_\_\_\_ 261 Aaron Stutz

\_\_\_\_\_ 271 Vanessa Sun

\_\_\_\_\_ 211 Jamey Kain

\_\_\_\_\_ 221 Laura Anderson

\_\_\_\_\_ 361 Reema Thalji

\_\_\_\_\_ 371 Warren Wood

\_\_\_\_\_ 311 David Barry

\_\_\_\_\_ 321 Christina Brown

\_\_\_\_\_ 461 Aaron Stutz

\_\_\_\_\_ 471 Alex Kollias

\_\_\_\_\_ 411 Andrew Chi

\_\_\_\_\_ 421 Shahed Ghoghawala

\_\_\_\_\_ 431 Mike Slater

\_\_\_\_\_ 561 Jean Han

\_\_\_\_\_ 571 Ognjen Miljanic

\_\_\_\_\_ 511 Joe Kwon

\_\_\_\_\_ 521 Catherine Chan

\_\_\_\_\_ 531 Olga Fedin

\_\_\_\_\_ 541 Jason Serin

If you are making up an I-grade, indicate the semester you took 3A  
\_\_\_\_\_ and the Professor \_\_\_\_\_

This exam has **15** pages; **make sure that you have them all.**

We will only grade answers that are in the designated spaces. Please do your scratch work on the backs of the exam pages. Write only **one** answer to each problem; multiple answers will receive **no** credit, even if one of them is correct.

**Note:** This examination runs for a total of 180 minutes. No questions will be answered by proctors after the exam begins. Please write legibly; ambiguous or messy answers will receive **no credit**.

© 2002 J.M.J. Fréchet, Berkeley, CA

**Do not write in this box.**

1. \_\_\_\_\_ (18)

2. \_\_\_\_\_ (15)

3. \_\_\_\_\_ (15)

4. \_\_\_\_\_ (11)

5. \_\_\_\_\_ (15)

6. \_\_\_\_\_ (14)

7. \_\_\_\_\_ (18)

8. \_\_\_\_\_ (12)

9. \_\_\_\_\_ (18)

10. \_\_\_\_\_ (15)

11. \_\_\_\_\_ (18)

12. \_\_\_\_\_ (14)

13. \_\_\_\_\_ (17)

**TOTAL** (1) (200)

Note: There are no questions to be answered on this page,  
Note that some of the data provided may not be needed

**Typical  $^1\text{H}$  NMR  
chemical shifts  $\delta$**

R-CH <sub>3</sub>	0.7-1.3
R-CH <sub>2</sub> -R'	1.2-1.6
R <sub>3</sub> CH	1.4-1.8
R-CH <sub>2</sub> -O-R'	3.3-4.0
R-CH <sub>2</sub> Cl	3.4-3.7
R <sub>2</sub> CHCl	3.9-4.4
RO-CH <sub>2</sub> Cl	5.2-5.6
RR'C=CH <sub>2</sub>	4.6-5.0
RCH=CHR'	5.2-5.7
RC≡CH	1.7-3.1
$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{R}$	2.0-2.4
$\text{C}=\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-$	1.7-2.2

**Typical  $^{13}\text{C}$  NMR  
chemical shifts  $\delta$**

R-CH <sub>3</sub>	5-25
R-CH <sub>2</sub> -R'	25-35
R <sub>3</sub> CH	35-55
R <sub>4</sub> C	30-45
R-CH <sub>2</sub> -O-R'	50-90
R-CH <sub>2</sub> Cl	25-50
R-COOH	170-180
R-CH=CH-R'	100-150
R <sub>2</sub> C=CR' <sub>2</sub>	100-150
RC≡CR	65-95
$\text{CH}_3-\overset{\text{O}}{\parallel}{\text{C}}-\text{R}$	170-220

**Infrared stretching in  $\text{cm}^{-1}$**

O-H (alcohol)	3200-3650
C-H (alkane)	2840-3000
C-H (alkene)	3050-3150
C-H (alkyne)	3260-3330
C=O (aldehyde, ketone)	1690-1750
C=C (alkene)	1620-1680
C≡C (alkyne)	2100-2260
$\text{R}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OR}'$ (ester)	1735-1750

D is the symbol used to denote deuterium  
the isotope  $^2\text{H}$  of hydrogen

**Bond dissociation energies (in Kcal mole<sup>-1</sup>):**


$\text{H}_2\text{C}=\overset{\cdot\cdot}{\text{C}}\text{H}-\overset{\cdot\cdot}{\text{H}}$	108 ✓	$\text{H}_2\text{C}=\overset{\cdot\cdot}{\text{C}}\text{H}-\overset{\cdot\cdot}{\text{Cl}}$	88 ✓
$\text{H}_2\text{C}=\overset{\cdot\cdot}{\text{C}}\text{H}_2$	both sigma and pi bonds: 153 pi bond only: 65.	$\text{RO}-\overset{\cdot\cdot}{\text{C}}\text{H}_2$	92 ✓
		$\text{RCH}_2-\overset{\cdot\cdot}{\text{C}}\text{H}-\overset{\cdot\cdot}{\text{Cl}}$	80 ✓
		$\text{RO}-\overset{\cdot\cdot}{\text{C}}\text{H}-\overset{\cdot\cdot}{\text{H}}$	104 ✓
		$\text{RO}-\overset{\cdot\cdot}{\text{C}}\text{H}-\overset{\cdot\cdot}{\text{Cl}}$	53 ✓

**Partial periodic table of the elements**



PERIOD	1					2		
1	<b>H</b> 1.008					<b>He</b> 4.003		
2	<b>3</b> <b>Li</b> 6.941	<b>4</b> <b>Be</b> 9.012	<b>5</b> <b>B</b> 10.81	<b>6</b> <b>C</b> 12.01	<b>7</b> <b>N</b> 14.01	<b>8</b> <b>O</b> 16.00	<b>9</b> <b>F</b> 19.00	<b>10</b> <b>Ne</b> 20.18
3	<b>11</b> <b>Na</b> 22.99	<b>12</b> <b>Mg</b> 24.31	<b>13</b> <b>Al</b> 26.98	<b>14</b> <b>Si</b> 28.09	<b>15</b> <b>P</b> 30.97	<b>16</b> <b>S</b> 32.06	<b>17</b> <b>Cl</b> 35.45	<b>18</b> <b>Ar</b> 39.95

1. (18 Points). Name or draw the following molecules as appropriate (show stereochemistry)


(a) (S)-4-heptyn-3-ol




(b)



(c)




(d)  $\text{H}-\text{C}\equiv\text{C}-\text{C}\equiv\text{C}-\text{CH}_3$




(e)

(E)-(2R,3R)-3-ethyl-4-hexen-2-ol













(f)




Cis-1,2-diethynylcyclohexane







2. (15 Points) Complete the following reactions showing the reagents used to effect the desired transformation. In some cases more than one step is required, in such cases, all reagents and the order in which they are used must be shown









(pure cis isomer)



3. (15 Points). Complete the following reactions showing the major reaction product(s). show stereochemistry when relevant. Write NR if no reaction takes place.





show most stable chair conformation of product!




4. (11 Points)

(a) An unknown compound  $C_4H_9Cl$  has the  $^1H$  NMR spectrum shown to the right. What is its structure?

Answer:




(b) The free radical chlorination of alkanes with sulfuryl chloride  $ClSO_2Cl$  does not produce the same mixture of monochlorinated products as would be obtained using  $Cl_2$  and light. Using the product compositions below, calculate the selectivity of the  $\cdot SO_2Cl$  radical for primary, and secondary hydrogens. Show the details of your work.




Answer. Relative reactivity **secondary/primary** =

5. (15 Points) (a) The IR spectrum below is for a linear alkyne with the molecular formula  $C_8H_{14}$ . Write a clear structure for this hydrocarbon.


Answer:





- (b) Consider the mechanism shown for the isomerization reaction below and draw a clear, fully labeled potential energy diagram showing clearly the relative location of the key species A, B, C, and of any transition states.



6. (14 Points). (a) Show a detailed step by step mechanism (with curved arrows) explaining the outcome of the following reaction:





- (b) Write the expected major product(s) of the following attempted ether syntheses







7. (18 Points). (a) Propose a reasonable structure for the organic compounds below that show only one peak in their  $^{13}\text{C}$  NMR spectra.

 $\text{C}_5\text{H}_{10}$  $\text{C}_4\text{H}_{12}\text{Si}$ 



(b) How many peaks are in the  $^{13}\text{C}$  NMR "normal" spectrum of:

Answer:



(c) How many peaks are in the DEPT-135  $^{13}\text{C}$  NMR spectrum of:

Answer:



(d) Give the approximate pKa values for the functional groups shown below

 $\text{R}-\text{C}\equiv\text{C}-\text{H}$  $\text{R}-\text{SH}$  $\text{R}-\text{OH}$ 


Answers:



(e) Which of the following bond lengths match most closely the lengths of the following carbon-carbon bonds (write *one* answer in each box below)?

Bond length in Angstroms (Å): 1.14 1.20 1.30 1.34 1.40 1.44 1.50 1.54

C-C

 $\text{C}\equiv\text{C}$ 

8. (12 Points) (a) A compound has the molecular formula  $C_7H_{12}O_2$ . What is its degree of unsaturation?

Answer:

(b) An unknown compound  $C_7H_{12}O_2$  has the following  $^{13}C$  NMR.

"Normal" proton decoupled spectrum,  $\delta$  values: 19, 28, 70, 129, 130, 166

DEPT-90 spectrum,  $\delta$  values: 28, 130

DEPT-135 spectrum,  $\delta$  values: *positive* peaks at 19, 28, 130; *negative* peaks at 70, 129

(i) What type(s) of protons are observed by DEPT-90? Answer:



(ii) What type(s) of protons give rise to negative peaks in DEPT-135?

Answer:

(iii) What is the structure of the unknown compound  $C_7H_{12}O_2$ ? **EXPLAIN** your reasoning.

Answer:

9. (18 Points) Consider the two reactions of ethene shown below:



each involving the initiation step:  $\text{RO}-\text{Cl} \xrightarrow{h\nu} \text{RO}\cdot + \text{Cl}\cdot$  [ $\Delta H^\circ = +53 \text{ Kcal mole}^{-1}$ ]

**Note** that for each reaction, the initiation step is followed by reaction of  $\text{RO}\cdot$  with ethene

(i) write the two propagation steps for the reaction (a) and calculate the  $\Delta H^\circ$  for each step

$$\Delta H^\circ =$$

$$\Delta H^\circ =$$





(ii) write the two propagation steps for the reaction (b) and calculate the  $\Delta H^\circ$  for each step

$$\Delta H^\circ =$$













$$\Delta H^\circ =$$

(iii) Explain why only one of the two reactions (**specify which one**) takes place at room temperature


10. (15 Points) Propose a step-by-step synthesis of  $\text{CH}_3\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}-\underset{\text{OH}}{\text{CH}}-\text{CH}_2\text{CH}_3$   
 from ethyne and propene as the sole sources of C atoms



11. (18 Points). Complete the following reactions showing the major product(s) or reagents. Show clear stereochemistry where relevant.



12. (14 Points) Propose a synthesis of 3-methylpentane from ethene used as the sole source of C atoms.




from  $\text{CH}_2=\text{CH}_2$

13. (17 Points) (a) Calculate the % of S enantiomer in a mixture of 2 enantiomers with an optical rotation of +12 degrees, given that the optical rotation of the pure S enantiomer is +15 degrees.

Answer:  %

(b) Give the IUPAC name of:








(c) What is the shape of the molecule of  $N(CH_3)_3$ ?

(d) Which is the strongest acid?

- (1)  $H-C\equiv C-H$  (2)  $NH_3$  (3)  $CH_4$  (4)  $H_2O$  (5)  $H_2C=CH-CH_3$  (6)  $CH_3COO^-$  (7)  $I^-$
- release H<sup>+</sup>, give*


Answer:

(e) Which of the following are meso compounds? Write the answer(s) in the box provided but do not guess as wrong answers will result in point deductions from correct answers.



Answer(s):

(f) Draw a Fischer projection of the product obtained in the reaction below and write the rate law for this reaction.



Answer:

Rate =