


Please solve all three problems below, and please explain your answers.

Problem 1 [30pts]: Spin-orbit interaction




The spin-orbit interaction $H'_{so} = \left(\frac{e^2}{8\pi\epsilon_0}\right) \frac{1}{m^2 c^2 r^3} \vec{S} \cdot \vec{L}$ causes the degenerate $2P$ level of the hydrogen atom (the level with principal quantum number $n = 2$ and orbital quantum number $l = 1$) to split into two levels: $2P_{\frac{1}{2}}$ (with total angular momentum quantum number $j = \frac{1}{2}$) and $2P_{\frac{3}{2}}$ (with total angular momentum quantum number $j = \frac{3}{2}$). They differ in energy by

$$\Delta E = E(2P_{\frac{3}{2}}) - E(2P_{\frac{1}{2}}) = 4.53 \times 10^{-5} \text{eV}.$$

Hypothetically, if the electron had spin $s = 1$ (instead of $s = \frac{1}{2}$), assuming the same form of H'_{so} , into how many distinct levels would $2P$ split, and what would be the energy differences between adjacent levels? (You can express your answers in terms of ΔE .)

Problem 2 [40pts]: Fermions and perturbation theory.

Two identical and non-interacting fermions of mass m are in an infinite square well potential in the range $-a < x < a$. We ignore the spin in this problem (assuming that both particles are in the same spin state).



- (a) What is the energy of the ground state of the system?
- (b) Write down the wavefunction $\psi(x_1, x_2)$ of the ground state of the system.
- (c) A perturbation $H' = \lambda x_1 x_2$ is added to the Hamiltonian. Calculate the first order $O(\lambda)$ correction to the ground state energy.

You can use one or more of the formulas

$$\int_{-a}^a \cos^2\left(\frac{\pi x}{2a}\right) dx = \int_{-a}^a \sin^2\left(\frac{\pi x}{a}\right) dx = a,$$

$$\int_{-a}^a x \cos\left(\frac{\pi x}{2a}\right) \sin\left(\frac{\pi x}{a}\right) dx = \frac{32a^2}{9\pi^2}, \quad \int_{-a}^a x \sin^2\left(\frac{\pi x}{a}\right) dx = \int_{-a}^a x \cos^2\left(\frac{\pi x}{2a}\right) dx = 0,$$

$$\int_{-a}^a x^2 \sin^2\left(\frac{\pi x}{a}\right) dx = \frac{a^3}{3} - \frac{a^3}{2\pi^2}, \quad \int_{-a}^a x^2 \cos^2\left(\frac{\pi x}{2a}\right) dx = \frac{a^3}{3} - \frac{2a^3}{\pi^2},$$

(some formulas were added to hide the really needed one.)

Please turn the page \implies

Problem 3 [30pts]: quick one-line answers.

Please answer the following questions and add to your answers one-line explanations.

- (a) Two spin- $\frac{3}{2}$ particles are in a state $|\frac{3}{2}, \frac{3}{2}\rangle$ where both z -components of spin have eigenvalues $+\frac{3}{2}\hbar$. What would a measurement of the square of the total spin $(\vec{S}_1 + \vec{S}_2)^2$ give?
- (b) Two identical fermions have an orbital wavefunction

$$\psi(\vec{r}_1, \vec{r}_2) = \frac{1}{\pi b^3} e^{-(r_1+r_2)/b}.$$

Considering the following possibilities for their spin-part of the wavefunction:

$$(1) \quad |\uparrow\uparrow\rangle; \quad (2) \quad |\uparrow\downarrow\rangle; \quad (3) \quad |\downarrow\uparrow\rangle; \quad (4) \quad |\downarrow\downarrow\rangle;$$

pick one of the following choices:

- (A) Exactly one answer from (1)-(4) is correct. [Which one is it?];
(B) there is more than one correct answer (1)-(4). [Which ones are correct?];
(C) none of the answers (1)-(4) is correct.

- (c) Taking into account fine-structure, one finds that the lowest 6 energy levels of the hydrogen atom are

$$E_1 = -13.60587\text{eV}, \quad E_2 = -3.401480\text{eV}, \quad E_3 = -3.401434\text{eV}, \\ E_4 = -1.511763\text{eV}, \quad E_5 = -1.511750\text{eV}, \quad E_6 = -1.511746\text{eV}$$

Write down the multiplicities (degeneracies) of each level. (Ignore the hyperfine splitting and the factor of 2 due to the spin of the proton.)

You may wish to recall the formula

$$E_{nj} = -\frac{1}{2n^2} mc^2 \alpha^2 + \frac{1}{n^4} \left(\frac{3}{4} - \frac{n}{j + \frac{1}{2}} \right) mc^2 \alpha^4, \quad \left(\alpha \equiv \frac{e^2}{4\pi\epsilon_0\hbar c} = \frac{1}{137.035999173(35)} \right)$$

Formulas for the midterm

Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{H}\psi, \quad \hat{H} = \frac{1}{2m}\hat{p}^2 + V(x), \quad \hat{p} = \frac{\hbar}{i} \frac{d}{dx}$$

Charged particle in a magnetic field

$$\hat{H} = \frac{1}{2m} \left(\frac{\hbar}{i} \vec{\nabla} - q\vec{A} \right)^2 - \vec{\mu} \cdot \vec{B}, \quad \vec{B} = \vec{\nabla} \times \vec{A}, \quad \vec{\mu} = \frac{gq}{2m} \vec{S},$$

Infinite square well

$$\begin{aligned} \psi_{2n+1} &= \frac{1}{\sqrt{a}} \cos\left(\frac{\pi x}{2a}\right), & E_{2n+1} &= \frac{\hbar^2(2n+1)^2}{8ma^2}, & (2n+1) &= 1, 3, 5, 7, \dots \\ \psi_{2n} &= \frac{1}{\sqrt{a}} \sin\left(\frac{\pi x}{a}\right), & E_{2n} &= \frac{\hbar^2(2n)^2}{8ma^2}, & (2n) &= 2, 4, 6, 8, \dots \end{aligned} \quad V(x) = \begin{cases} 0, & \text{if } |x| > a \\ \infty, & \text{if } -a < x < a \end{cases}$$

Harmonic oscillator

$$\begin{aligned} \hat{H} &= \frac{1}{2m}\hat{p}^2 + \frac{1}{2}m\omega^2\hat{x}^2 = \hbar\omega\left(a_+a_- + \frac{1}{2}\right), & [a_-, a_+] &= 1, & \hat{a}_{\pm} &= \frac{1}{\sqrt{2\hbar m\omega}}(\mp i\hat{p} + m\omega\hat{x}), \\ \psi_0(x) &= \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-\frac{m\omega}{2\hbar}x^2}, & \psi_n &= \frac{1}{\sqrt{n!}}(a_+)^n\psi_0, & E_n &= (n + \frac{1}{2})\hbar\omega \end{aligned}$$

Spherical harmonics

$$\begin{aligned} P_l(x) &= \frac{1}{2^l l!} \left(\frac{d}{dx}\right)^l [(x^2 - 1)^l], & P_l^m(x) &= (1 - x^2)^{|m|/2} \left(\frac{d}{dx}\right)^{|m|} P_l(x), \\ Y_{lm}(\theta, \phi) &= \epsilon \sqrt{\frac{(2l+1)}{4\pi} \frac{(l-|m|)!}{(l+|m|)!}} e^{im\phi} P_l^m(\cos\theta), & \epsilon &= \begin{cases} (-1)^m & \text{for } m \geq 0 \\ 1 & \text{for } m < 0 \end{cases} \\ Y_{0,0} &= \frac{1}{\sqrt{4\pi}}, & Y_{1,0} &= \left(\frac{3}{4\pi}\right)^{1/2} \cos\theta, & Y_{1,\pm 1} &= \mp \left(\frac{3}{8\pi}\right)^{1/2} \sin\theta e^{\mp i\phi}, \\ Y_{2,0} &= \left(\frac{5}{16\pi}\right)^{1/2} (3\cos^2\theta - 1), & Y_{2,\pm 1} &= \mp \left(\frac{15}{8\pi}\right)^{1/2} \sin\theta \cos\theta e^{\mp i\phi}, & Y_{2,\pm 2} &= \left(\frac{15}{32\pi}\right)^{1/2} \sin^2\theta e^{\mp 2i\phi}, \end{aligned}$$

Angular momentum

$$\begin{aligned} \hat{L}_{\pm} &\equiv L_x \pm iL_y = \pm \hbar e^{\pm i\phi} \left(\frac{\partial}{\partial \theta} \pm i \cot\theta \frac{\partial}{\partial \phi} \right), & \hat{L}^2 &= -\hbar^2 \left[\frac{1}{\sin\theta} \frac{\partial}{\partial \theta} \left(\sin\theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2\theta} \frac{\partial^2}{\partial \phi^2} \right] \\ \hat{L}_z |l, m\rangle &= \hbar m |l, m\rangle, & \hat{L}^2 |l, m\rangle &= \hbar^2 l(l+1) |l, m\rangle, & \hat{L}_{\pm} |l, m\rangle &= \hbar \sqrt{l(l+1) - m(m \pm 1)} |l, m \pm 1\rangle \end{aligned}$$

Pauli matrices

$$\begin{aligned} \sigma_x &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, & \sigma_y &= \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, & \sigma_z &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & \vec{S} &= \frac{\hbar}{2} \vec{\sigma}, \\ \sigma_x^2 &= \sigma_y^2 = \sigma_z^2 = \mathbf{I}, & [\sigma_x, \sigma_y] &= 2i\sigma_z, & [\sigma_y, \sigma_z] &= 2i\sigma_x, & [\sigma_z, \sigma_x] &= 2i\sigma_y, \end{aligned}$$

Hydrogen-like Atom

$$\psi_{100} = \frac{1}{\sqrt{\pi a^3}} e^{-r/a}, \quad a = \frac{4\pi\epsilon_0 \hbar^2}{me^2} = 0.529 \times 10^{-10} \text{m}, \quad E_1 = -\frac{m}{2\hbar^2} \left(\frac{e^2}{4\pi\epsilon_0} \right)^2 = -13.6 \text{eV}, \quad E_n = \frac{E_1}{n^2}$$

$$\psi_{nlm} = R_{nl}(r) Y_{lm}(\theta, \phi), \quad R_{10} = \frac{2}{\sqrt{a^3}} e^{-x}, \quad R_{20} = \frac{1}{\sqrt{2a^3}} (1 - \frac{1}{2}x) e^{-x/2}, \quad R_{21} = \frac{1}{\sqrt{24a^3}} x e^{-x/2}, \quad x \equiv \frac{r}{a}$$

Identical particles

$$\psi(x_1, x_2) = \begin{cases} \frac{1}{\sqrt{2}} [\psi_1(x_1)\psi_2(x_2) - \psi_2(x_1)\psi_1(x_2)] & \text{noninteracting identical fermions} \\ \psi_1(x_1)\psi_1(x_2) \quad \text{or} \quad \frac{1}{\sqrt{2}} [\psi_1(x_1)\psi_2(x_2) + \psi_2(x_1)\psi_1(x_2)] & \text{noninteracting identical bosons} \end{cases}$$

singlet: $\frac{1}{2}(|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$, triplet: $|\uparrow\uparrow\rangle, \frac{1}{2}(|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle), |\downarrow\downarrow\rangle$.

Time-independent perturbation theory

$$E_n^{(1)} = \langle \psi_n^{(0)} | H' | \psi_n^{(0)} \rangle, \quad \psi_n^{(1)} = \sum_{m \neq n} \frac{\langle \psi_m^{(0)} | H' | \psi_n^{(0)} \rangle}{E_n^{(0)} - E_m^{(0)}} \psi_m^{(0)}, \quad E_n^{(2)} = \sum_{m \neq n} \frac{|\langle \psi_m^{(0)} | H' | \psi_n^{(0)} \rangle|^2}{E_n^{(0)} - E_m^{(0)}} \psi_m^{(0)}$$

Degenerate perturbation theory

$$W_{ij} \equiv \langle \psi_i^{(0)} | H' | \psi_j^{(0)} \rangle, \quad E_{\pm}^{(1)} = \frac{1}{2} \left[W_{aa} + W_{bb} \pm \sqrt{(W_{aa} - W_{bb})^2 + 4|W_{ab}|^2} \right]$$

Fine structure

$$H' = -\frac{\hat{p}^4}{8m^3c^2} + \left(\frac{e^2}{4\pi\epsilon_0} \right) \frac{1}{m^2c^2r^3} \vec{S} \cdot \vec{L}, \quad \vec{S} \cdot \vec{L} = \frac{1}{2}(J^2 - L^2 - S^2), \quad \vec{J} = \vec{L} + \vec{S}.$$

$$E_{nj} = -\frac{\alpha^2 mc^2}{2n^2} \left[1 + \frac{\alpha^2}{n^2} \left(\frac{n}{j+\frac{1}{2}} - \frac{3}{4} \right) \right], \quad \alpha \equiv \frac{e^2}{4\pi\epsilon_0 \hbar c} = \frac{1}{137.035999173(35)}$$