

EECS 70  
Fall 2015

Discrete Mathematics and Probability Theory  
Jean Walrand

Midterm 3

PRINT Your Name: \_\_\_\_\_,  
(last) (first)

SIGN Your Name: \_\_\_\_\_

PRINT Your Student ID: \_\_\_\_\_

CIRCLE your exam room: 2040 VLSB 2060 VLSB 145 Dwinelle 155 Dwinelle 10 Evans OTHER

Name of the person sitting to your left: \_\_\_\_\_

Name of the person sitting to your right: \_\_\_\_\_

- After the exam starts, please write your student ID (or name) on every odd page (we will remove the staple when scanning your exam).
- We will not grade anything outside of the space provided for a problem. Please use scratch paper as necessary and clearly indicate your answer.
- For questions 1(a)-(e). You need only circle True or False.
- For questions 2 (a)-(g), only provide the requested answer (e.g., probability value, one or more events). There is no need to justify your answer.
- For questions 3 (a)-(d), write clearly your answer in the space provided. There is no need to justify your answer.
- For questions 4 (a)-(h), you should indicate clearly your derivation in the space provided.
- You may not look at books, notes, etc. Calculators, phones, and computers are not permitted.
- There are 9 pages on the exam, including this first page. Notify a proctor immediately if a page is missing.
- **You may, without proof, use theorems and facts that were proven in the notes and/or in lecture.**
- **You have 105 minutes; there are 24 parts on this exam.**

Do not turn this page until your instructor tells you to do so.

**1. True or False. No justification needed. 15 points. 3/3/3/3.**

**Clearly indicate your correctly formatted answer: this is what is to be graded. No need to justify!**

- (a) Disjoint events with a positive probability cannot be independent. (True or False.)
  
- (b) We can find events  $A$  and  $B$  with  $Pr[A|B] > Pr[A]$  and  $Pr[B|A] < Pr[B]$ . (True or False.)
  
- (c) If  $Pr[A|B] = Pr[B]$ , then  $A$  and  $B$  are independent. (True or False.)
  
- (d) For a random variable  $X$ , it is always the case that  $E[X^2 - X] \geq -1$ . (True or False)
  
- (e) If  $Pr[A] > Pr[\bar{A}]$ , then  $Pr[A|B] \geq Pr[\bar{A}|B]$ . (True or False)

**2. Short Answer: Probability Space. 31 points: 4/4/4/5/5/4/5**

**Clearly indicate your correctly formatted answer: this is what is to be graded. No need to justify!**

- (a) You flip a biased coin (such that  $Pr[H] = p$ ) until you accumulate two  $H$ s (not necessarily consecutive). What is the probability space? That is, what is  $\Omega$  and what is  $Pr[\omega]$  for each  $\omega \in \Omega$ ?

SID:

---

(b) Let  $\Omega = \{1, 2, 3, 4\}$  be a uniform probability space. Let also  $A = \{1, 2, 3\}$ . Produce an event  $B$  such that  $Pr[B] > 0$  and  $A$  and  $B$  are independent.

(c) Let  $\Omega = \{1, 2, 3, 4\}$  be a uniform probability space. Produce three events  $A, B, C$  that are pairwise independent but not mutually independent.

(d) You are dealt two cards from a deck of 52 cards. What is the probability that the value of the first card is strictly larger than that of the second? [In this question, the values are 1 for an ace, 2 through 10 for the number cards, then 11 for a Jack, 12 for a queen, 13 for a king.]

SID:

---

(e) You roll a balanced 6-sided die twice. What is the probability that the total number of pips is less than 10 given that it is larger than 7?

(f) With probability  $1/2$ , one rolls a die with four equally likely outcomes  $\{1, 2, 3, 4\}$  and with probability  $1/2$  one rolls a balanced die with six equally likely outcomes  $\{1, 2, \dots, 6\}$ . Given that the outcome is 4, what is the likelihood that the coin was four-sided?

(g) A coin is equally likely to be fair or such that  $Pr[H] = 0.6$ . You flip the coin 10 times and get 10 heads. What is the probability that the next coin flip yields heads?



**4. Short Problems. 40 points: 5/5/5/5/5/5****Clearly indicate your answer and your derivation.**

(a) Let  $X$  be a random variable with mean 1. Show that  $E[2 + 3X + 3X^2] \geq 8$ .

(b) Let  $X$  be geometrically distributed with parameter  $p$ . Recall that this means that  $Pr[X = n] = (1 - p)^{n-1}p$  for  $n \geq 1$ . Find  $E[X|X > n]$ . Do not leave the answer as an infinite sum.

SID:

---

(c) Roll a die  $n$  times. Let  $X_n$  be the average number of pips per roll. What is  $\text{var}[X_n]$ ? You may leave the answer as a sum.

(d) Let  $X$  and  $Y$  be independent with  $X = G(p)$  and  $Y = G(q)$ . What is  $\Pr[X \leq Y]$ ? Do not leave the answer as an infinite sum.

SID:

---

(e) You roll a balanced die five times. Let  $X$  be the total number of pips you got and  $Y$  the total number of pips on the last two rolls. What is  $E[X|Y = 4]$ ? What is  $E[Y|X = 15]$ ?

(f) How many times do you have to flip a fair coin, on average, until you get two consecutive  $H$ 's? [Hint: condition on the outcome of the last flip.]



- (g) Let  $\{X_n, n \geq 1\}$  be independent and geometrically distributed with parameter  $p$ . Recall that  $\text{var}[X] = (1-p)/p^2$ . Provide an upper bound on

$$\Pr\left[\left|\frac{X_1 + \cdots + X_n}{n} - p\right| \geq a\right]$$

using Chebyshev's inequality.

- (h) There are two envelopes. One contains checks with  $\{1, 3, 5, 6, 7\}$  dollars. The other contains checks with  $\{4, 5, 5, 7\}$  dollars. You choose one of the two envelopes at random and pick one of the checks at random in the envelope. That check happens to be for 5 dollars. You are given the option to keep all the money in that envelope, including the check for 5 dollars, or to switch to the other envelope. What should you do?