STUDENT ID:

Math 54, first midterm, Fall 2018, John Lott

This is a closed book, closed notes, closed calculator, closed computer, closed phone, closed network, open mind exam.

Be sure to write your name and student id number on the top of EVERY page that you turn in.

Name of neighbor to your left	
Name of neighbor to your right _	

Write your answers in the boxes provided. For full credit, show your work, too, (except for the true/false questions) and cross out work that you do not want us to grade.

You can write on the backs of pages. You can also use the back of this page, and the back of the last page, as scratch paper. If you need additional scratch paper, ask me for it.

STUDENT ID:

1. (10 pts) Is $\begin{bmatrix} 3 \\ -7 \\ -3 \end{bmatrix}$ in the span of the vectors

$$\begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}, \begin{bmatrix} -4 \\ 3 \\ 8 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ -4 \end{bmatrix}?$$

Write YES or NO in the box.



Justify your answer.

NO

The augmented matrix is

$$\begin{bmatrix} 1 & -4 & 2 & 3 \\ 0 & 3 & 5 & -7 \\ -2 & 8 & -4 & -3 \end{bmatrix}.$$

Performing a row operation gives

$$\begin{bmatrix} 1 & -4 & 2 & 3 \\ 0 & 3 & 5 & -7 \\ 0 & 0 & 0 & 3 \end{bmatrix}.$$

The last row is inconsistent.

2. (10 pts) (No partial credit, but show your work.)

Write the inverse of

$$\begin{bmatrix} 1 & 2 & 4 \\ 1 & 3 & 5 \\ 1 & 2 & 3 \end{bmatrix}$$

in the box.

The answer is

$$\begin{bmatrix} 1 & -2 & 2 \\ -2 & 1 & 1 \\ 1 & 0 & -1 \end{bmatrix}.$$

STUDENT ID:

3. (10 pts) A linear transformation T from \mathbb{R}^2 to \mathbb{R}^3 satisfies

$$T\left(\begin{bmatrix}1\\1\end{bmatrix}\right) = \begin{bmatrix}4\\0\\0\end{bmatrix}, \quad T\left(\begin{bmatrix}1\\-1\end{bmatrix}\right) = \begin{bmatrix}0\\6\\0\end{bmatrix}.$$

Write the standard matrix of T in the box. Justify your answer.



$$\begin{bmatrix} 2 & 2 \\ 3 & -3 \\ 0 & 0 \end{bmatrix}.$$

This is because

$$T\left(\begin{bmatrix}1\\0\end{bmatrix}\right) = \begin{bmatrix}2\\3\\0\end{bmatrix}, \quad T\left(\begin{bmatrix}0\\1\end{bmatrix}\right) = \begin{bmatrix}2\\-3\\0\end{bmatrix}.$$

STUDENT ID:

4. (12 pts) Suppose that B is an invertible square matrix with the property that for both B and B^{-1} , all of their entries are integers. Show that $\det(B)$ is 1 or -1.

$$\det(B)\det(B^{-1}) = \det(I) = 1.$$

Since B and B^{-1} both have integer entries, det(B) and $det(B^{-1})$ are both integers, so det(B) must be ± 1 .

STUDENT ID:

WARNING: there are		No explanation is necessary. s problem. If you copy off of a neighbor, you wrong.
If $\{\mathbf{x_1}, \mathbf{x_2}, \mathbf{x_3}, \mathbf{x_4}\}$ is independent.	a set of linearly independ	dent vectors in \mathbb{R}^4 then $\{\mathbf{x_1}, \mathbf{x_2}, \mathbf{x_3}\}$ is linearly
	\square TRUE	\square FALSE
If A is an $m \times n$ matrix	atrix then the null space	of A is a subspace of \mathbb{R}^m .
	☐ TRUE	☐ FALSE
If A is a matrix and	$A^5 = I$ then A is inver	tible.
If A and B are $n \times$	n matrices, and AB is in	nvertible, then A and B are invertible. \Box FALSE
If a matrix A has line	nearly dependent columi	as then $A\mathbf{x} = 0$ has infinitely many solutions. \Box FALSE

TA T		7	/		
17	Δ	- 11	∕∎	н,	•

STUDENT ID:

NAME:		IUDENI ID:			
The set of vectors $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$	so that $x_1 + x_2 + x_3$ $\square \text{ TRUE}$	that $x_1 + x_2 + x_3 \ge 0$ forms a subspace of \mathbb{R}^3 . \square TRUE \square FALSE			
If A is a 5×5 matrix	such that $dot(2.4)$ —	dot(A) then $A=0$			
If A is a 5×5 matrix	such that $det(2A) =$	$\det(A)$ then $A=0$.			
	\square TRUE	\square FALSE			
If T is a one-to-one li	near transformation t	from \mathbb{R}^n to \mathbb{R}^n then T is onto.			
II I IS a One-to-one in					
	\square TRUE	\square FALSE			
If V is a vector space the set of vectors that li		subspaces, then the union of H_1 and vays a subspace.	H_2 (i.e.		
	\square TRUE	\square <u>FALSE</u>			