

FINAL EXAMINATION  
Chemistry 3A

Name: Key SID #: \_\_\_\_\_  
Print first name before second! *Make sure the number is correct!*  
Use capital letters!

GSI (if you are taking Chem 3AL): \_\_\_\_\_

Peter Vollhardt  
May 15, 2019

**Please write the last four digits of your Cal ID# on the back of each page of this exam.**

Provide the following information if applicable.

Making up an I Grade \_\_\_\_\_

If you are, please indicate the semester during which you took previous Chem 3A and the instructor:

\_\_\_\_\_      \_\_\_\_\_  
Semester      Instructor

Auditor \_\_\_\_\_

**Please write the answer you wish to be graded in the boxed spaces provided.**

**Do not remove this (or any other) page from the exam. It will compromise the conformance of scanned copies in Gradescope.**

**Do scratch work on the back of the pages.** This test should have **25** numbered pages. Check to make sure that you have received a complete exam. A good piece of advice: **Read carefully over the questions (at least twice); make sure that you understand exactly what is being asked; avoid sloppy structures or phrases. It is better to be pedantic in accuracy now than sorry later! Good Luck!**

You probably will not need it for the exam, but here it is for added comfort:

Electronegativity values of the elements (Pauling scale)

H 2.1																	He
Li 1.0	Be 1.5											B 2.0	C 2.5	N 3.0	O 3.5	F 4.0	Ne
Na 0.9	Mg 1.2											Al 1.5	Si 1.8	P 2.1	S 2.5	Cl 3.0	Ar
K 0.8	Ca 1.0	Sc 1.3	Ti 1.5	V 1.6	Cr 1.6	Mn 1.5	Fe 1.8	Co 1.8	Ni 1.8	Cu 1.9	Zn 1.6	Ga 1.6	Ge 1.8	As 2.0	Se 2.4	Br 2.8	Kr 3.0
Rb 0.8	Sr 1.0	Y 1.2	Zr 1.4	Nb 1.6	Mo 1.8	Tc 1.9	Ru 2.2	Rh 2.2	Pd 2.2	Ag 1.9	Cd 1.7	In 1.7	Sn 1.8	Sb 1.9	Te 2.1	I 2.5	Xe 2.6
Cs 0.7	Ba 0.9	La 1.1	Hf 1.3	Ta 1.5	W 1.7	Re 1.9	Os 2.2	Ir 2.2	Pt 2.2	Au 2.4	Hg 1.9	Tl 1.8	Pb 1.8	Bi 1.9	Po 2.0	At 2.2	Rn 2.4
Fr 0.7	Ra 0.7	Ac 1.1															
Ce 1.1	Pr 1.1	Nd 1.1	Pm 1.1	Sm 1.1	Eu 1.1	Gd 1.1	Tb 1.1	Dy 1.1	Ho 1.1	Er 1.1	Tm 1.1	Yb 1.1	Lu 1.2				
Th 1.3	Pa 1.5	U 1.7	Np 1.3	Pu 1.3	Am 1.3	Cm 1.3	Bk 1.3	Cf 1.3	Es 1.3	Fm 1.3	Md 1.3	No 1.3	Lr				

**Table 2.2** Relative Acidities of Common Compounds (25°C)


Acid	$K_a$	$pK_a$
Hydrogen iodide, HI (strongest acid)	$\sim 1.0 \times 10^{10}$	-10.0
Hydrogen bromide, HBr	$\sim 1.0 \times 10^9$	-9.0
Hydrogen chloride, HCl	$\sim 1.0 \times 10^8$	-8.0
Sulfuric acid, $H_2SO_4$	$\sim 1.0 \times 10^3$	-3.0 <sup>a</sup>
Hydronium ion, $H_3O^+$	50	-1.7
Nitric acid, $HNO_3$	25	-1.4
Methanesulfonic acid, $CH_3SO_3H$	16	-1.2
Hydrogen fluoride, HF	$6.3 \times 10^{-4}$	3.2
Acetic acid, $CH_3COOH$	$2.0 \times 10^{-5}$	4.7
Hydrogen cyanide, HCN	$6.3 \times 10^{-10}$	9.2
Ammonium ion, $NH_4^+$	$5.7 \times 10^{-10}$	9.3
Methanethiol, $CH_3SH$	$1.0 \times 10^{-10}$	10.0
Methanol, $CH_3OH$	$3.2 \times 10^{-16}$	15.5
Water, $H_2O$	$2.0 \times 10^{-16}$	15.7
Ethyne, $HC\equiv CH$	$\sim 1.0 \times 10^{-25}$	$\sim 25$
Ammonia, $NH_3$	$1.0 \times 10^{-35}$	35
Ethene, $H_2C=CH_2$	$\sim 1.0 \times 10^{-44}$	$\sim 44$
Methane, $CH_4$ (weakest acid)	$\sim 1.0 \times 10^{-50}$	$\sim 50$

Note:  $K_a = [H_3O^+][A^-]/[HA]$  mol L<sup>-1</sup>.

<sup>a</sup>First dissociation equilibrium

- I. [60 Points] Provide the IUPAC name or draw the structure, as appropriate, of the following molecules. Remember the priority of functional groups in choosing names, indicate the correct stereochemistry (e.g. *R*, *S*, *cis*, *trans*, and *E*, *Z*), and do not forget about the alphabetical ordering of substituents!

a.




This enantiomer

(*R,E*)-5-Ethynyl-4-(prop-2-ynyl)octa-4,7-dien-3-ol


b.

(*R,Z*)-2-(2-Chloro-2-butenyl)octane-1-thiol



Complete the stencil

c.




(This is a Fischer projection;  
name this enantiomer)

(*S*)-3-Methylpent-1-en-4-yne

d.


Meso-1,2-dimethoxy-1,2-bis(methylthio)cyclohexane



Complete the stencil


e.

(*R*)-3-Methylpent-1-en-4-yne



Complete the Fischer stencil.




f.



*Cis*-(*E*)-1-Iodo-4-(1,1,2-trimethylbut-2-enyl)-cyclohexane


II. [100 Points] Add the missing starting materials, reagents, or products (aqueous work-up is assumed where necessary). Don't forget **stereochemistry**! Complete the stencils, when provided. **Note:** In the reagent boxes above the arrows, enter one step only. Multistep sequences are clearly indicated by several boxes labelled "1., 2., 3.," etc.

a.




Two diastereomers

b.



This enantiomer



For the following questions, circle your choice of an answer:

Is the product chiral?

Yes



No

Is the product optically active?



Yes

No




c.




d.




e.

1.  $\text{Br}_2, h\nu$ 2.  $\text{NaOCH}_3, \text{CH}_3\text{OH}$ 3.  $\text{O}_3$ 4.  $(\text{CH}_3)_2\text{S}$ 


f.




g.




h.



i.




j.





III. [45 Points] Examine the potential radical addition of ethane to 1-propene to give 2-methylbutane (Eq. 1).



a. Using  $DH^\circ_{\text{C}-\text{H}} = 65 \text{ kcal mol}^{-1}$  and the appropriate bond dissociation data in the Table below, calculate the  $\Delta H^\circ$  of Eq. 1. Show your work.

**Selected C–C Bond-Dissociation Energies (kcal mol<sup>-1</sup>)**

$\text{CH}_3-\dot{\text{C}}\text{H}_3$	90
$\text{C}_2\text{H}_5-\dot{\text{C}}\text{H}_3$	89
$\text{CH}_3\text{CH}_2\text{CH}_2-\dot{\text{C}}\text{H}_3$	89
$\text{C}_2\text{H}_5-\dot{\text{C}}\text{H}_2$	88
$(\text{CH}_3)_2\text{CH}-\dot{\text{C}}\text{H}_3$	88
$(\text{CH}_3)(\text{CH}_3\text{CH}_2)\text{CH}-\dot{\text{C}}\text{H}_3$	88
$(\text{CH}_3)_3\text{C}-\dot{\text{C}}\text{H}_3$	87
$(\text{CH}_3)_2\text{CH}-\dot{\text{C}}\text{H}(\text{CH}_3)_2$	85.5
$(\text{CH}_3)_3\text{C}-\dot{\text{C}}(\text{CH}_3)_3$	78.5

$\Delta H^\circ$  of Eq. 1:

$$(65 + 90) - (88 + 89) = -22 \text{ kcal mol}^{-1}$$


b. Considering that the entropy of the reaction is negative and makes an unfavorable contribution to its  $\Delta G^\circ$ , amounting to  $\sim +10 \text{ kcal mol}^{-1}$  (at 25°C), is the reaction thermodynamically (**circle** your answer):

Favorable?

Unfavorable?


c. Write the two initiation steps for the radical chain reaction and calculate their respective  $\Delta H^\circ$  values, using an  $\text{CH}_3\text{O}-\text{OCH}_3$  bond strength of  $39 \text{ kcal mol}^{-1}$  and  $DH^\circ(\text{CH}_3-\text{O}) = 90 \text{ kcal mol}^{-1}$ .

Step 1.



$$\Delta H^\circ = +39 \text{ kcal mol}^{-1}$$


Step 2.



$$\Delta H^\circ = 0 \text{ kcal mol}^{-1}$$


e. Write the two propagation steps for the reaction in Eq. 1 and calculate their respective  $\Delta H^\circ$  values using the data given in part (a).

Step 1.



$$\Delta H^\circ = -24 \text{ kcal mol}^{-1}$$

Step 2.




$$\Delta H^\circ = 2 \text{ kcal mol}^{-1}$$

f. Which propagation step would you consider to be relatively slow?  
(Place an "X" mark next to your answer in the box provided).

Step 1


Step 2

IV. [20 Points] Compound **A** enters into two equilibria. The first is with **B**, proceeds with an activation energy of 40 kcal mol<sup>-1</sup>, and is favorable with  $\Delta G^\circ = -30$  kcal mol<sup>-1</sup>. The second is with **C**, proceeds with an activation energy of 30 kcal mol<sup>-1</sup>, and is favorable with  $\Delta G^\circ = -10$  kcal mol<sup>-1</sup>.



Complete the energy diagram below as follows:

- Starting with **A** positioned at level "0" (black bar), indicate the respective positions of **B** (to the right) and **C** (to the left) along the reaction coordinate (use black bars and label them with the respective letters).
- Show the positions of the two respective transition states (black bars) that reflect the activation energies of the two steps. Label them "**TS<sub>AB</sub>**" and "**TS<sub>AC</sub>**", respectively.



V. [20 Points] Using key words or a short sentence, list three observations that support the mechanism of the E1 reaction.

1.

Rate =  $k[R-L]$ : unimolecular kinetics

2.


Stereochemistry: racemization

3.


Rate increases with increasingly better leaving group

VI. [75 Points] Write detailed step-wise mechanisms for the following two transformations (a. and b.) Use only structures and “arrow-pushing of electron dots” techniques. Note: These are not synthetic problems. Do not add any reagents! What you see is what you have!


a.




Work from left to right in the following spaces. There is much more space than you will need.



b.




Work from left to right in the following spaces. There is much more space than you will need. Do not forget to include all electron dots.




c. On treatment with base, Compound **A** undergoes intramolecular Williamson ring closure faster than its diastereomer **B**. Why?

1. To find the answer, draw the two chair conformers of **A** and **B** using the stencils in the boxes below. Add H, CH<sub>3</sub>, Br, or OH to the ends of the dangling bonds.



$$\Delta G^\circ = -0.21 \text{ kcal mol}^{-1}$$



$$\Delta G^\circ = +3.19 \text{ kcal mol}^{-1}$$

2. Using the table below, calculate the  $\Delta G^\circ$  values for the respective ring flips depicted in the boxes provided above.

**Table 4.3** Change in Free Energy on Flipping from the Cyclohexane Conformer with the Indicated Substituent Equatorial to the Conformer with the Substituent Axial

Substituent	$\Delta G^\circ$ [kcal mol <sup>-1</sup> (kJ mol <sup>-1</sup> )]	Substituent	$\Delta G^\circ$ [kcal mol <sup>-1</sup> (kJ mol <sup>-1</sup> )]
H	0 (0)	F	0.25 (1.05)
CH <sub>3</sub>	1.70 (7.11)	Cl	0.52 (2.18)
CH <sub>3</sub> CH <sub>2</sub>	1.75 (7.32)	Br	0.55 (2.30)
(CH <sub>3</sub> ) <sub>2</sub> CH	2.20 (9.20)	I	0.46 (1.92)
(CH <sub>3</sub> ) <sub>3</sub> C	≈ 5 (21)	HO	0.94 (3.93)
	1.41 (5.90)	CH <sub>3</sub> O	0.75 (3.14)
	1.29 (5.40)	H <sub>2</sub> N	1.4 (5.9)





3. Armed with the results of the above analysis, why does **A** react faster than **B**? Explain in one sentence.

Unlike **A**, **B** needs to flip to a higher-energy conformer to adopt the anti arrangement of Br and OH necessary for the intramolecular  $S_N2$  reaction.

VII. [20 Points] The following reactions proceed (predominantly) by  $S_N2$ ,  $S_N1$ , E2, or E1 pathways, respectively. Give the major products in each case and answer the questions by **circling** the most applicable statement.

a.



Mechanism:

$S_N2$

$S_N1$

**E2**

E1

When using  $\text{Na}^+\text{SH}$  instead of  $\text{Na}^+\text{OC}(\text{CH}_3)_3$ , which one of the following ratios will increase:



$S_N2 / S_N1$

$S_N1 / E1$

$E2 / S_N2$

**$S_N2 / E2$**

b.



Mechanism:

**$S_N2$**

$S_N1$

E2

E1

When changing the  $-\text{N}(\text{CH}_3)_2$  group to  $-\text{P}(\text{CH}_3)_2$ , the rate of disappearance of starting material:


**Will increase**

Will decrease


Will not change

VIII. [90 Points] Show synthetic **forward connections** (reagents, intermediates; no mechanisms!) between the following starting materials and the final products. Note: several steps are required in each case; there may be several solutions to each problem, but you should show only one. It is best to work backwards (retrosynthetically) on the back of the exam pages, to enable you to dissect the products into less complex precursors. However, the answer to be graded should be a *forward* scheme.


a. Starting from **A** and any organic and organometallic reagents **containing four carbons or less**, propose a synthesis of **B**.








Work from left to right in the following spaces. There is more space than you will need.




b. Provide a scheme using **only the starting material** to supply the necessary building blocks for assembling the product.




Work from left to right in the following spaces.



c.





Work from left to right in the following spaces. There is much more space than you will need.



## IX. [70 Points]


a. Which of the following potential energy diagrams describes the reaction depicted below left? Give your answer (**A**, **B**, **C**, or **D**) in the box provided.



Answer:

**D**

b. Rank the following alkenes in the order of increasing rates of oxacyclopropanation:



Place an "X" mark next to your answer.


\_\_\_ A, B, C, E, D

\_\_\_ D, B, A, E, C

\_\_\_ B, C, D, E, A

\_\_X\_\_ A, D, E, B, C

c. Solvolysis of **B** is one million times slower than that of **A**. Why?



Place an "X" mark next to your answer.


\_\_\_ **B** is more sterically hindered.

\_\_\_ The electron pairs on fluorine are electron donating.

\_\_\_ **B** is better solvated.


\_\_**X**\_\_  $\text{CF}_3$  is electron withdrawing, destabilizing the tertiary cation.

d. Draw the most important contributing resonance form of the species shown below:



e. The molecule shown below has primary, secondary, and tertiary hydrogens. Considering the relative reactivity in radical brominations of C–H: primary : secondary : tertiary = 1 : 80 : 1700, what would you expect the ratio of **all** primary to **all** secondary to **all** tertiary substitution products to be?


Show your work and just give the raw ratio.



primary : secondary : tertiary = 1 : 80 : 1700

$1 \times 12 : 80 \times 12 : 1700 \times 2 = 12 : 960 : 3400$

f. Consider the bromohydroxylation ( $\text{Br}_2, \text{H}_2\text{O}$ ) of the alkene enantiomer shown below. How many bromoalcohol isomers do you expect?




(*R,Z*)-4-Methyl-2-heptene

Place an "X" mark next to your answer.

- two enantiomers  
 four diastereomers  
 two chiral and two meso compounds  
 four isomers  
 eight stereoisomers


g. Consider the following substitution reaction:



Suggest one experiment that would help you decide whether **this particular reaction** occurs by  $\text{S}_{\text{N}}2$  or  $\text{S}_{\text{N}}1$ .

Double  $[\text{H}_2\text{S}]$ :  
 if rate doubles =  $\text{S}_{\text{N}}2$   
 if rate is unchanged =  $\text{S}_{\text{N}}1$





\* The End \*

*Have a great summer!*