

Chem 4A Exam 1

TOTAL POINTS

94 / 100

QUESTION 1

1 1A 10 / 10

- ✓ + **4 pts** Correct balanced chemical reaction
- ✓ + **2 pts** Correct ideal gas law
- ✓ + **0.5 pts** Correct mole fraction
- ✓ + **3 pts** Correct final answer
- ✓ + **0.5 pts** Correct units

QUESTION 2

2 1B 10 / 10

- ✓ + **1 pts** Use definition of molarity
- ✓ + **2 pts** Correct moles of propanol
- ✓ + **1 pts** Convert to moles propanol to CO₂
- ✓ + **2 pts** Equation/mole conversion is balanced
- ✓ + **1 pts** Convert moles CO₂ to grams
- ✓ + **1 pts** Correctly use molar mass of CO₂
- ✓ + **1 pts** Correct significant figures
- ✓ + **1 pts** Correct units
- **1 pts** Math error
- + **0 pts** Incorrect

QUESTION 3

3 2A 10 / 10

- ✓ + **5 pts** Correct final answer
- ✓ + **3.5 pts** Correct half-reaction
- ✓ + **1.5 pts** Evidence of valid stoichiometry (attempts are counted even with wrong numbers as long as dimensional analysis is valid so as to not double count for mistakes)
- + **2.25 pts** Correct thought process for obtaining half reaction, but incorrect equation (cannot coincide with "correct half-reaction")
- + **4.5 pts** Answer off by a reasonable factor (in that I can track your mistake) + almost correct stoichiometry (cannot coincide with "correct final answer" or

"evidence of valid stoichiometry"); this does not count if your half reaction is wrong

- + **1 pts** Correct dimensional analysis without the right numbers (cannot coincide with "evidence of valid stoichiometry"), this is a special case so as not to double count for mistakes or double count for correctness
- + **4.5 pts** Correct final answer but wrong number of sigfigs
- + **0 pts** Incorrect and incomplete

QUESTION 4

4 2B 10 / 10

- ✓ + **2 pts** Have the correct equation set up
- ✓ + **5 pts** correct calculation to answer
- ✓ + **3 pts** Get the right answer (36mL) from part A
- **1 pts** Sig figs
- + **0 pts** incorrect

QUESTION 5

5 3A 10 / 10

- ✓ + **0.5 pts** Mass to mol conversions for Cu
- ✓ + **2 pts** Conservation equations
- ✓ + **4 pts** Solution to the conservation equation
- ✓ + **2 pts** Mass of each component
- ✓ + **1 pts** Sig fig
- ✓ + **0.5 pts** Mass to mol conversion for O
- + **0 pts** Incorrect

QUESTION 6

6 3B 10 / 10

- ✓ + **2 pts** Correct mean
- ✓ + **2 pts** Correct standard deviation
- ✓ + **2 pts** Correct t and N
- ✓ + **4 pts** Correct confidence interval based on previously calculated values

+ 0 pts Incorrect

QUESTION 7

7 SO₃ 2- 10 / 10

- ✓ + 2 pts Best Overall Structure
- ✓ + 2 pts Correct Formal Charge
- ✓ + 1 pts Correct Lone Pairs
- ✓ + 1 pts Correct Geometry
- ✓ + 2 pts Correct Electron Pair Geometry
- ✓ + 2 pts Correct Molecular Geometry
- + 0 pts Wrong molecule
- + 0 pts Incorrect

✓ + 2 pts Structure

✓ + 1 pts Electrons

✓ + 1 pts Geometry

+ 0 pts Incorrect

QUESTION 8

8 NO⁺ 5 / 10

- 0 pts Correct
- ✓ - 2 pts Formal Charge
- ✓ - 1 pts Lone Pair
- 1 pts Geometry
- ✓ - 2 pts Structure
- 1 pts Half Correct Structure
- 2 pts Electron Geometry
- 2 pts Molecular Geometry

QUESTION 9

9 NO₂ 9 / 10

- ✓ + 2 pts formal charge
- ✓ + 2 pts electron pair geometry
- ✓ + 2 pts molecular geometry
- ✓ + 2 pts Structure
- ✓ + 1 pts electrons
- ✓ + 1 pts Geometry
- + 0 pts [Click here to replace this description.](#)
- + 0 pts [Click here to replace this description.](#)
- 1 Point adjustment
- ☹ Not best structure

QUESTION 10

10 PO₄ 3- 10 / 10

- ✓ + 2 pts Formal charge
- ✓ + 2 pts Electron pair geometry
- ✓ + 2 pts Molecular geometry

Chemistry 4A, Exam I
September 16, 2019
Professor R. J. Saykally

N _____

1. (20) _____
2. (20) _____
3. (20) _____
4. (40) _____

TOTAL EXAM SCORE (100) _____

Rules:

- Work all problems to 2 significant figures
- No lecture notes or books permitted
- No programmable or graphing calculators permitted
- Time: 50 minutes
- Show all work to get partial credit
- All answers must be written in the boxes provided
- Periodic Table, Tables of Physical Constants, and Conversion Factors included

Physical Constants


Standard Acceleration of terrestrial gravity	$g = 9.80665 \text{ m s}^{-2}$ (exactly)
Avogadro's number	$N_0 = 6.022137 \times 10^{23}$
Bohr radius	$a_0 = 0.52917725 \text{ \AA} = 5.2917725 \times 10^{-11} \text{ m}$
Boltzmann's constant	$k_B = 1.38066 \times 10^{-23} \text{ J K}^{-1}$
Electron Charge	$e = 1.6021773 \times 10^{-19} \text{ C}$
Faraday constant	$\mathcal{F} = 96,485.31 \text{ C mol}^{-1}$
Masses of fundamental particles:	
Electron	$m_e = 9.109390 \times 10^{-31} \text{ kg}$
Proton	$m_p = 1.672623 \times 10^{-27} \text{ kg}$
Neutron	$m_n = 1.674929 \times 10^{-27} \text{ kg}$
Ratio of proton mass to electron mass	$m_p/m_e = 1836.15270$
Permittivity of vacuum	$\epsilon_0 = 8.8541878 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$
Planck's constant	$h = 6.626076 \times 10^{-34} \text{ J s}$
Speed of light in vacuum	$c = 2.99792458 \times 10^8 \text{ m s}^{-1}$ (exactly)
Universal gas Constant	$R = 8.31451 \text{ J mol}^{-1} \text{ K}^{-1} = 0.0820578 \text{ L atm mol}^{-1} \text{ K}^{-1}$
Rydberg Constant	$R_\infty = e^4 m_e / (8 \epsilon_0^2 h^2)$

Conversion Factors


Standard Atmosphere	$1 \text{ atm} = 1.01325 \times 10^5 \text{ Pa} = 1.01325 \times 10^5 \text{ kg m}^{-1} \text{ s}^{-2}$ (exactly)
Atomic mass unit	$1 \text{ u} = 1.660540 \times 10^{-27} \text{ kg}$
	$1 \text{ u} = 1.492419 \times 10^{-10} \text{ J} = 931.4942 \text{ MeV}$ (energy equivalent from $E = mc^2$)
Calorie	$1 \text{ cal} = 4.184 \text{ J}$ (exactly)
Electron volt	$1 \text{ eV} = 1.6021773 \times 10^{-19} \text{ J} = 96.48531 \text{ kJ mol}^{-1}$
Foot	$1 \text{ ft} = 12 \text{ in} = 0.3048 \text{ m}$ (exactly)
Gallon (U.S.)	$1 \text{ gallon} = 4 \text{ quarts} = 3.78541 \text{ L}$ (exactly)
Liter-atmosphere	$1 \text{ L atm} = 101.325 \text{ J}$ (exactly)
Metric ton	$1 \text{ metric ton} = 1000 \text{ kg}$ (exactly)
Pound	$1 \text{ lb} = 16 \text{ oz} = 0.45359237 \text{ kg}$ (exactly)

Question 1 (10 points each)

A) What mass (g) of CO_2 is made when 2.0 L of a stoichiometric mixture of gaseous propanol ($\text{C}_3\text{H}_7\text{OH}$) and oxygen at 1.0 atm, 300 K is combusted?




B) Calculate the mass of CO_2 produced by complete surface combustion of 0.70 L of a 4.0 M solution of propanol ($\text{C}_3\text{H}_7\text{OH}$) in water.



Question 2 (10 points each)


The Iodine Clock Reaction involves the reaction of iodate (IO_3^-) with iodide (I^-) in acidic solution (H^+) to produce iodine (I_2) and water.

- A) Calculate the volume of 0.100 M IO_3^- solution that will exactly react with 20.0 mL of 0.300 M I^- solution.



- B) What volume of 0.100 M sulfite (SO_3^{2-}) solution would be required to exactly react with the iodine (I_2) produced in Part A above?

The balanced reaction is: $\text{SO}_3^{2-} + \text{I}_2 + \text{H}_2\text{O} \rightarrow \text{SO}_4^{2-} + 2\text{I}^- + 2\text{H}^+$



Question 3 (10 points each)

A. ✓ 1.000-g mixture of cuprous oxide, Cu_2O , and cupric oxide, CuO , was quantitatively reduced to 0.839 g of metallic copper. What was the mass of CuO in the original sample?

$$x = \text{g Cu}_2\text{O} \quad y = \text{g CuO} \quad 2\text{Cu}_2\text{O} \rightarrow 4\text{Cu} + \text{O}_2 \quad 2\text{CuO} \rightarrow 2\text{Cu} + \text{O}_2$$

$$x + y = 1.000 \text{ g} \quad \text{Cu} = 63.55 \text{ g/mol} \quad \text{Cu}_2\text{O} = \frac{143.099 \text{ g}}{\text{mol}} \quad \text{CuO} = \frac{79.544 \text{ g}}{\text{mol}}$$

$$x = 1.000 - y$$

$$\frac{x \text{ g Cu}_2\text{O}}{143.099 \text{ g/mol}} \cdot \frac{4 \text{ mol Cu}}{2 \text{ mol Cu}_2\text{O}} \cdot \frac{63.55 \text{ g Cu}}{\text{mol Cu}} = 0.8882x \text{ g Cu}$$

$$\frac{y \text{ g CuO}}{79.544 \text{ g/mol}} \cdot \frac{2 \text{ mol Cu}}{2 \text{ mol CuO}} \cdot \frac{63.55 \text{ g Cu}}{\text{mol Cu}} = 0.7989y \text{ g Cu}$$

$$0.8882x + 0.7989y = 0.839$$

$$0.8882(1.000 - y) + 0.7989y = 0.839$$

$$0.0893y = 0.0492$$

$$y = \frac{0.0492}{0.0893} \approx 0.55 \text{ g CuO}$$

$$\boxed{0.55 \text{ g CuO}}$$

B) Four trials yield the following results for the mass of CuO produced in the above reactions:

0.538 g 0.716 g 0.815 g 0.920 g

Calculate the 95% confidence interval for these results.

Table 1: "t" Values for 95% confidence interval.

Degrees of freedom	Value of "t"
1	6.314
2	2.920
3	2.353
4	2.132
5	2.015
6	1.943

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n} = \frac{0.538 + 0.716 + 0.815 + 0.920}{4} = 0.747 \text{ g}$$

$$s_x = \sqrt{\frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n-1}} = \sqrt{\frac{(0.538 - 0.747)^2 + (0.716 - 0.747)^2 + (0.815 - 0.747)^2 + (0.920 - 0.747)^2}{4-1}} = 0.162$$

$$s_x = 0.162 \text{ g}$$

$$\bar{x} \pm \frac{t s}{\sqrt{n}}$$

$$t: \text{Dof} = 4 - 1 = 3 \rightarrow t = 2.353$$

$$\bar{x} \pm \frac{2.353 \cdot 0.162}{\sqrt{4}} = 0.747 \text{ g} \pm 0.191 \text{ g}$$

↓ 2 sig figs


$$\boxed{0.75 \text{ g} \pm 0.19 \text{ g}}$$

Question 4 (10 points each)


For the following compounds:

- Draw the Lewis Structure, explicitly showing the formal charges and molecular geometry
- Indicate the Electron Pair Geometry
- Indicate the Molecular Geometry

I. Sulfite Ion (SO_3^{2-})

A. 	B. tetrahedral	C. trigonal pyramidal
---	-------------------	--------------------------

II. Nitronium Ion (NO^+)

A. 	B. Linear	C. Linear
---	--------------	--------------

III. ✓

NO₂ Molecule
SN=3

A. 	B. <i>trigonal planar</i>	C. <i>bent</i>
--------	----------------------------------	-------------------

IV. ✓

Phosphate Ion (PO₄³⁻)
SN=4

A. 	B. <i>tetrahedral</i>	C. <i>tetrahedral</i>
--------	--------------------------	--------------------------